This is a digital copy of a book that was preserved for generations on library shelves before it was carefully scanned by Google as part of a project to make the world's books discoverable online.

It has survived long enough for the copyright to expire and the book to enter the public domain. A public domain book is one that was never subject to copyright or whose legal copyright term has expired. Whether a book is in the public domain may vary country to country. Public domain books are our gateways to the past, representing a wealth of history, culture and knowledge that's often difficult to discover.

Marks, notations and other marginalia present in the original volume will appear in this file - a reminder of this book's long journey from the publisher to a library and finally to you.

Usage guidelines

Google is proud to partner with libraries to digitize public domain materials and make them widely accessible. Public domain books belong to the public and we are merely their custodians. Nevertheless, this work is expensive, so in order to keep providing this resource, we have taken steps to prevent abuse by commercial parties, including placing technical restrictions on automated querying.

We also ask that you:

+ *Make non-commercial use of the files* We designed Google Book Search for use by individuals, and we request that you use these files for personal, non-commercial purposes.

+ *Refrain from automated querying* Do not send automated queries of any sort to Google's system: If you are conducting research on machine translation, optical character recognition or other areas where access to a large amount of text is helpful, please contact us. We encourage the use of public domain materials for these purposes and may be able to help.

+ *Maintain attribution* The Google "watermark" you see on each file is essential for informing people about this project and helping them find additional materials through Google Book Search. Please do not remove it.

+ *Keep it legal* Whatever your use, remember that you are responsible for ensuring that what you are doing is legal. Do not assume that just because we believe a book is in the public domain for users in the United States, that the work is also in the public domain for users in other countries. Whether a book is still in copyright varies from country to country, and we can't offer guidance on whether any specific use of any specific book is allowed. Please do not assume that a book's appearance in Google Book Search means it can be used in any manner anywhere in the world. Copyright infringement liability can be quite severe.

About Google Book Search

Google's mission is to organize the world's information and to make it universally accessible and useful. Google Book Search helps readers discover the world's books while helping authors and publishers reach new audiences. You can search through the full text of this book on the web at http://books.google.com/

QL95
F77

J. P.

HARVARD UNIVERSITY
ARNOLD ARBORETUM
PRESENTED BY CHARLES SPRAGUE SARGENT
1892

FLORVLAE

INSVLARVM AVSTRALIVM

PRODROMVS

AVCTORE

GEORGIO FORSTER M. D.

SERENISSIMO REGI POLONIAE A CONSILIIS INTIMIS HISTORIAE NATVRALIS ET BOTANICES IN ACADEMIA VILNENSI P. P. O. ACADEMIAE CAESAREAE NATVRAE CVRIOSORVM REGIAE MEDICAE MADRITENSIS REGIAR. SOCIETATVM SCIENTIARVM LONDINENSIS ET HAVNIENSIS SOCIETATVM ANTIQVITATVM ET REI AGRARIAE QVAE CASSELLIS SVNT REI AGRARIAE CELLENSIS ET NATVRAE SCRVTATOR. BEROLINENSIS SODALI NEC NON REGIIS SOCIETATIB. GOTTINGENSI SCIENTIARVM ET PARISINAE REI AGRARIAE LITTERARVM COMMERCIO CONIVNCTO.

GOTTINGAE

TYPIS JOANN. CHRISTIAN. DIETERICH.

MDCCLXXXVI.

VIRO

EXCELLENTISSIMO ET CELEBERRIMO

ANDREAE SPARRMANNO

MEDICINAE DOCTORI
PROFESSORI HISTORIAE NATVRALIS
MVSEI HOLMIENSIS ACADEMICI PRAEFECTO
ACADEMIAE REGIAE SCIENTIARVM HOLMIENSIS
SOCIETATIS PHYSIOGRAPHICAE LVNDINENSIS
MEMBRO &c.

FLORVLAE

COMMVNI STVDIO ORNATAE

RECENSVM

D.

GEORGIVS FORSTER.

PRAEFATIO.

Ex itinere circa globum terraqueum redux anno MDCCLXXV, coniunctim cum patre optimo *Characterum nouorum generum plantarum* adumbrationem typis tabellisque aeneis exaratam publici iuris feci. Hanc breui, vt ſperabam, ita pollicebar deſcriptiones integras ſtirpium orbis auſtralis inſequuturas. Haud igitur ab re eſſet enumerare cauſſas, quae harum diuulgationem impediverunt; verum piget querelis quamuis iuſtis immorari.

Interim vndecimus ab editione nouorum generum vertitur annus. Species plantarum complures, a nobis primum detectae, alienis curis prodierunt in lucem, ſiue quod in ſolo natali iterum deprehenſae fuerunt, ſiue quod, etſi a nobis mutuatae, alieno nomini minus religioſe illae addictae ſunt. Facile quidem tolerari poterat iactura tantillae, inuenti legitimis auctoribus erepti, gloriolae. Sed non potuit non vitium ſubrepere deſcriptiunculis alienis iuxta ſicca plantarum exemplaria congeſtis et feſtinanter nimis in medium prolatis. Quae cum ita eſſent, iam vltro videbar adſtipulari erroribus, ni publice in eosdem animaduerterem. Attamen eadem, quae olim, obſtacula manum noſtro operi admotam inhibent. Qui in colligendas haſce naturae diuitias aes publicum impenderunt, ad earundem communicationem cum orbe erudito ſumtus denegant, librarii redemtionem recuſant; vetat res familiaris

familiaris tot figurarum ſcalpturas, quae deſcriptionibus illuſtrandis inſeruiunt, priuatis opibus moliri. Quid multa? Praemittendum itaque exploratoris vice catalogum Florulae Polyneſiae conſtitui, certiorem me redditurum, num in promtu mihi ſint auxiliatrices manus, quibus omnibus his anguſtiis expediar? Enimuero, niſi cuncta me fallunt, praeſagiens iam animo tot inter potentiſſimos Europae principes exorientem Muſagetam video, qui noſtrorum tentaminum fautorem ſe praeſtet, Floraeque auſtralis ornandae honorem Britannis eripiat.

Nomina plantarum, quas cum amiciſſimo Sparrmanno in oceani auſtralis inſulis legimus deſcripſimusque, ſolis differentiis ſpecificis et locis natalibus adiectis, huic prodromo inſerui, excluſis ſtirpibus magellanicis, nec non iis quas in atlantici maris inſulis offendimus. Species cognitas omnes

ad

ad Illuſtris MURRAYI editionem decimam quartam ſyſtematis vegetabilium retuli, emendatis hinc inde differentiis ſpecificis, quas fide ſupplementi plantarum a LINNEO, filio, editi, in eximium opus ſuum receperat vir celeberrimus. Genera noua nonnulla, inſtituta diligentiore comparatione, deleui; quae autem temere nimis ab aliis ad genera cognita relata fuerunt, reſtitui. Plantas dubias porro et obſcuras aut imperfecte deſcriptas ſeorſim enumerandas eſſe iudicaui, ne et ſpecies probe determinatae vna vacillarent.

Scribebam Vilnae Lituanicae, d. XXX. Junii MDCCLXXXVI.

FLORVLAE

FLORVLAE
INSVLARVM AVSTRALIVM
PRODROMVS.

CLASSIS I.

MONANDRIA.

MONOGYNIA.

AMOMVM.

1. A. ***Zerumbet***, ſcapo nudo, ſpica oblonga obtuſa. MVRRAY *Syſtema Vegetabilium* edit. XIV. p. 50. n. 2.
Taheiti et reliquae Societatis inſulae.

2. A. ***Curcuma***, ſcapo ſpathis laxis e centro foliorum. M. *S. V.* p. 40. n. 5.
Culta intra tropicos et in Paſchatis inſula.

THALIA.

3. T. *cannaeformis*, corollis hexapetalis: nectario bifido erecto. F.
Mallicollo inter nouas Hebrides.

BOERHAAVIA.

4. B. *erecta*, caule erecto glabro, floribus diandris. M. *S. V.* p. 52. n. 1.
Inſulae Societatis. Forst. *pl. Eſculent. inſ. Oc. auſtr.* 41.

5. B. *tetrandra*, caule repente, floribus tetrandris. F. Ibidem.

DIGYNIA,

MNIARUM.

6. M. *biflorum*. M. *S. V.* p. 53. Forst. *Charact. Gen.* 1. Noua Zeelandia.

CLASSIS

CLASSIS II.

DIANDRIA.

MONOGYNIA.

IASMINVM.

7. I. *ſimplicifolium*, foliis oppoſitis ouato-lanceolatis ſimplicibus. F.
Amicorum Inſulae.

8. I. *didymum*, foliis oppoſitis ternatis, foliolis ouato-lanceolatis acuminatis integerrimis. F.
Societatis Inſulae.

VERONICA.

9. V. *Catarractae*, racemis terminalibus flexuoſis, caule ſuffruticoſo, foliis lanceolatis ſerratis F.
Noua Zeelandia.

10. V. *elliptica*, racemis lateralibus, caule fruticoſo, foliis ellipticis integerrimis. F.
Ibidem.

11. V. *ſalicifolia*, racemis lateralibus nutantibus, caule fruticoſo, foliis longo-lanceolatis integerrimis. F.
Ibidem.

IVSTICIA.

12. I. *repanda*, fruticoſa, foliis ouatis repandis: pedunculis axillaribus trifidis. F.
Tanna.

13. I. *longifolia*, ſuffruticoſa, foliis linearibus ſinuatis: pedunculis axillaribus trifidis. F.
Ibidem.

DIANTHERA.

14. D. *caerulea*, vmbella faſciculata ſeſſili ſimpliciſſima. F.
Botanices inſula, prope Nouam Caledoniam.

15. D. *clauata*, panicula ſubumbellata compoſita, pedunculis apice dilatatis. F.
Societatis inſulae.

GRATIOLA.

16. G. *Monnieria*, foliis ouali-oblongis, pedunculis vnifloris, caule repente. M. *S. V.* p. 64. n. 2.
Inſulae Marchionis.

VERBENA.

17. V. *nodiflora*, tetrandra, ſpicis capitato-conicis, foliis ſerratis, caule repente. M. *S. V.* p. 66. n. 10.
Tanna.

DIGYNIA.

ANTHOXANTHVM.

18. A. *crinitum*, panicula coarctata, ariſtis longiſſimis. F.
A. crinitum. M. *S. V.* p. 73. n. 4.
Noua Zeelandia.

TRIGYNIA.

PIPER.

19. P. *Siriboa*, foliis inaequaliter cordatis ſubſeptemneruiis venoſis. M. *S. V.* p. 74. n. 5.
Noua Caledonia.

20.

20. P. *excelſum*, foliis orbiculato-cordatis ſubſeptemneruiis: pedunculis terminalibus ſolitariis bifidis, caule arboreo. F.
Noua Zeelandia.

21. P. *methyſticum*, foliis cordatis acuminatis multineruiis: ſpicis axillaribus ſolitariis breuiſſimis pedunculatis patentiſſimis. FORST. *Plant. Eſc. Inſular. Oc. Auſtr.* 50.
Inſulae Societatis, Amicorum, Sandwigiae. Culta.

22. P. *latifolium*, foliis orbiculato-cordatis nouemnerviis: ſpicis axillaribus aggregatis pedunculatis. F.
P. latifolium. M. *S. V.* p. 74. n. 15.
Societatis, Amicorum, Nouarumque Hebridum inſulae.

23. P. *acuminatum?* foliis lanceolato-ouatis neruoſis carnoſis. M. *S. V.* p. 74. n. 11.
Societatis inſulae.

24. P. *pallidum*, foliis alternis obouatis ſubtrineruiis: ſpicis ſolitariis ſubterminalibus. F.
Ibidem.

25. P. *tetraphyllum*, foliis quaternis rhombeis ſeſſilibus carnoſis. F.
Ibidem.

CLASSIS III.

TRIANDRIA.

MONOGYNIA.

COMMELINA.

26. C. *virginica,?* corollis ſubaequalibus, foliis lanceolatis ore barbatis, caulibus erectis? M. *S. V.* p. 94. n. 5.
Tongatabu.

CYPERVS.

27. C. *odoratus*, culmo triquetro nudo, vmbella decompoſita ſimpliciter folioſa, pedicellis diſtiche ſpicatis. M. *S. V.* p. 97. n. 19.
Societatis inſulae.

SCIRPVS.

28. S. *dichotomus*, culmo triquetro nudo, vmbella decompoſita, ſpicis dichotomiae ſeſſilibus. M. *S. V.* p. 100. n. 24.
Societatis inſulae.

29. S. *junceus*, culmo teretiuſculo filiformi nudo, ſpica ouata terminali, glumis inuolucri longitudine ſpicae, tribus mucronatis. F.
Ibidem.

KYLLINGA.

30. K. *monocephala*, culmo filiformi triquetro, capitulo globoſo ſeſſili, inuolucro triphyllo longiſſimo. M. *S. V.* p. 101. n. 1.
Societatis inſulae.

31.

31. K. *triceps*, capitulis terminalibus ſubternis, glomeratis ſeſſilibus. M. *S. V.* p. 101. n. 2.
Thryocephalon nemorale Forst. *Charact. gen.* 65.
Ibidem.

DIGYNIA.

SACCHARVM.

32. S. *ſpontaneum*, panicula laxa, ſpicis filiformibus ſimplicibus, floribus remotis geminatis ariſtatis: altero pedunculato. F.
S. ſpontaneum M. *S. V.* p. 103. n. 1.
Societatis inſulae.

33. S. *officinarum*, floribus paniculatis, lanugine floſculis longiore. F. *pl. eſc.* 51.
S. officinarum. M. *S. V.* p. 103. n. 2.
Culta intra tropicos et in Paſchatis inſula.

PHALARIS.

34. P. *canarienſis*, panicula ſubouata ſpiciformi, glumis carinatis. M. *S. V.* p. 103. n. 1.
Noua Zeelandia.

PASPALVM.

35. P. *orbiculare*, ſpicis alternis, rachi membranacea, floribus duplici ordine ſecundis orbiculatis laeuibus. F.
Societatis inſulae.

PANICVM.

36. P. *hiſpidum*, ſpicis alternis ſubdiuiſis ſecundis, floſculis geminatis alternis ſecundis hiſpidis. F.
Noua Caledonia.

37\. P. *ſanguinale*, ſpicis digitatis baſi interiore nodoſis, floſculis geminis muticis, vaginis foliorum punctatis. M. *S. V.* p. 106. n. 13.
Societatis inſulae.

38\. P. *filiforme*, ſpicis ſubdigitatis approximatis erectis linearibus, rachi flexuoſa, dentibus bifloris: altero ſeſſili. M. *S. V.* p. 106. n. 15.
Ibidem et Paſchatis inſula.

39\. P. *compoſitum*, ſpica compoſita, ſpiculis linearibus ſecundis, floſculis geminis remotis, calycibus ariſtatis. M. *S. V.* p. 106. n. 18.
Ibidem.

AGROSTIS.

40\. A. *ouata*, petalo exteriore ante apicem ariſtato, panicula ouata, coarctata ſpiciformi. F.
Noua Zeelandia.

AIRA.

41\. A. *antarctica*, foliis planis, panicula compoſita patente, calycibus trifloris, floſculis medio ariſtatis, ariſta elongata rectiuſcula. F.
Noua Zeelandia.

POA.

42\. P. *Eragroſtis*, panicula patente, pedicellis flexuoſis, ſpiculis ſerratis decemfloris, glumis trineruiis. M. *S. V.* p. 114. n. 17.
Noua Caledonia.

43\. P. *anceps*, panicula diffuſa, ſpiculis quinquefloris acutis baſi pubeſcentibus, culmo ancipite. F.
Noua Zeelandia.

44\. P. *latifolia*, panicula diffuſa nutante, ſpiculis remotis bifloris, altero pedicellato, margine ſetoſo. F.

An

An Holcus latifolius? M. *S. V.* p. 905. n. 12.
Inſulae Societatis.

CYNOSVRVS.

45. C. *indicus*, ſpicis digitatis linearibus, culmo compreſſo declinato baſi nodoſo, foliis alternis. M. *S. V.* p. 117. n. 10.
Inſulae Societatis.

AVENA.

46. A. *filiformis*, panicula erecta tenuiſſima, calycibus vnifloris, ariſtis calyce duplo longioribus. F.
Noua Zeelandia et inſula Paſchatis.

ARVNDO.

47. A. *Bambos*, calycibus multifloris, ſpicis ternis ſeſſilibus. M. *S. V.* p. 123. n. 1.
Paſſim intra tropicos.

48. A. *conſpicua*, calycibus vnifloris, panicula laxa erecto-patente, petali exterioris ariſta reflexa longiſſima. F.
Noua Zeelandia.

ROTTBÖLLA.

49. R. *Coelorachis*, ſpica tereti vnilaterali, floſculis geminis altero pedicellato: calyce biualui. F.
Tanna.

50. R. *repens*, ſpica tereti ſubulata, gluma calycina vniualui indiuiſa. F.
Inſulae intra tropicos.

CLASSIS IV.

TETRANDRIA.

MONOGYNIA.

PETESIA.

51. P. *carnea*, foliis oblongo - lanceolatis laeuibus: cymis terminalibus trifidis. F.
Namoka.

ANCISTRVM.

52. A. *diandrum*, floribus diandris. F.
A. anſerinaefolium. Forst. *Charact. gen.* 2.
A. Sanguiſorbae. M. *S. V.* p. 73. n. 1.
Noua Zeelandia.

BLACKBVRNIA.

53. B. *pinnata.* Forst. *Charact. gen.* 6.
Norfolciae inſula deſerta.

FAGARA.

54. F. *Evodia*, foliis ternatis, foliolis integris. F.
F. Evodia. M. *S. V.* p. 160. n. 1.
Evodia hortenſis. Forst. *Charact. gen.* 7.
Inſulae Amicorum et nouae Hebrides.

OLDENLANDIA.

55. O. *foetida*, vmbella terminali trichotoma, foliis ſpatulatis. F.
Tonga - tabu.

56. O. *debilis*, vmbellis axillaribus pedunculatis pauci floris: foliis ouatis ſeſſilibus. F.
Ibidem.

57.

57. O. *tenuifolia*, pedunculis axillaribus ſolitariis vnifloris: foliis lineari-ſubulatis. F.
O. tenuifolia. BURM. *ind.* p. 47. tab. XIV. f. 1.
Tanna.

DORSTENIA.

58. D. *lucida*, caulescens, foliis oblique ouatis integris laeuibus, pedunculis cymoſis axillaribus. F.
Elatoſtema pedunculatum. FORST. *Charact. gen.* 53. n. 1.
Societatis inſulae.

59. D. *pubeſcens*, caulescens, foliis oblique ouatis ſerratis pubeſcentibus: pedunculis axillaribus capituliferis. F.
Elatoſtema ſeſſile. FORST. *Charact. gen.* 53. n. 2.
Ibidem.

EMBOTHRIVM.

60. E. *vmbellatum*, vmbellis axillaribus ſimpliciſſimis pedunculatis, antheris ſeſſilibus. FORST. *Charact. gen.* 8. n. 2.
E. vmbellatum. M. *S. V.* p. 166. n. 1.
Noua Caledonia.

TETRAGYNIA.

TILLAEA.

61. T. *muſcoſa?* procumbens, floribus trifidis. M. *S. V.* p. 170. n. 4.
Noua Zeelandia.

CLASSIS

CLASSIS V.

PENTANDRIA.

MONOGYNIA.

MYOSOTIS.

62. M. *ſpatulata*, ſeminibus laeuibus, foliis ſpatulatis hiſpidis, pedunculis axillaribus ſolitariis vnifloris. F.
Noua Zeelandia.

LITHOSPERMVM.

63. L. *incanum*, ſeminibus aſperis: ſpicis terminalibus compoſitis coarctatis, foliis linearibus villoſis. F. ♄.
Inſula Teautea et inſula ferox (Savage I).

TOURNEFORTIA.

64. T. *argentea*, foliis ouatis obtuſis tomentoſo-ſericeis: ſpicis terminalibus compoſitis. M. *S. V.* p. 191. n. 7.
Amicorum inſulae.

LYSIMACHIA.

65. L. *decurrens*, racemis ſimplicibus terminalibus, corollae ſegmentis obtuſis: ſtaminibus corolla longioribus. F.
Tanna.

OPHIORRHIZA.

66. O. *ſubumbellata*, caule fruticoſo, foliis lanceolatis acutis: vmbellis axillaribus trifidis. F.
Taheiti.

SHEFFIELDIA.

SHEFFIELDIA.

67. S. *repens*. Forst. *Charact. gen.* 9. M. *S. V.* p. 198. n. 1.
Noua Zeelandia et inſula Paſchatis.

EPACRIS.

* fructu capſulari.

68. E. *longifolia*, arborea, foliis lineari-lanceolatis baſi vaginantibus: racemis erectis articulatis lateralibus. F.
E. longifolia. Forst. *Charact. gen.* 10. n. 1. M. *S. V.* p. 198. n. 1.
Noua Zeelandia.

69. E. *roſmarinifolia*, fruticoſa, foliis linearibus obtuſis rigidis vaginantibus: floribus ſolitariis lateralibus. F.
Ibidem.

70. E. *pumila*, herbacea? foliis ouato-oblongis imbricatis: floribus ſeſſilibus ſubſolitariis terminalibus. F.
E. pumila. Forst. *Charact. gen.* 10. n. 3. M. *S. V.* p. 198. n. 3.
Ibidem.

** fructu baccato; *Stipheliae*, Solandr.

71. E. *juniperina*, arborea, foliis ſparſis linearibus cuſpidatis, ſerrulatis: floribus ſeſſilibus ſolitariis terminalibus. F.
E. juniperina. Forst. *Charact. gen.* 10. n. 2.
Ibidem.

72. E. *faſciculata*, arborea, foliis confertis linearibus acuminatis: racemis ſpicatis lateralibus cernuis. F.
E. juniperina. M. *S. V.* p. 198. n. 2.
Ibidem.

PLVMBAGO.

PLVMBAGO.

73. P. *zeylanica*, foliis petiolatis ouatis glabris, caulibus filiformibus. M. *S. V.* p. 199. n. 2.
Societatis inſulae.

CONVOLVVLVS.

74. C. *Tuguriorum*, (volubilis) foliis cordato-ſagittatis acutis: caule anguloſo, pedunculis tetragonis vnifloris. F.
Noua Zeelandia.

75. C. *Turpethum*, foliis cordatis angulatis, caule membranaceo-quadrangulari, pedunculis multifloris. M. *S. V.* p. 201. n. 22. FORST. *pl. Eſc.* 52.
Societatis et Amicorum inſulae.

76. C. *grandiflorus?* (volubilis) foliis cordatis acutis rugoſiuſculis multineruiis: calycis foliolis ſubrotundis obtuſiſſimis, pedunculis clauatis vnifloris. F.
C. grandiflorus? M. *S. V.* p. 201. n. 23.
Tanna.

77. C. *coeleſtis*, (volubilis) foliis cordatis acuminatiſſimis pubeſcentibus: pedunculis elongatis vmbellato-trifidis. F.
Tanna.

78. C. *peltatus*, foliis peltatis, pedunculis multifloris. M. *S. V.* p. 201. n. 27.
Societatis inſulae.

79. C. *mucronatus*, (volubilis) foliis palmato-pedatis, lobis ciliatis apice mucronatis: pedunculis vnifloris. F.
Tanna.

80. C. *corymboſus*, foliis cordatis, pedunculis vmbellatis, caule repente. M. *S. V.* p. 203. n. 52.
Noua Caledonia.

81.

81. C. *braſilienſis*, foliis emarginatis baſi biglanduloſis: pedunculis trifloris. M. *S. V.* p. 204. n. 62.
Intra tropicos paſſim.

IPOMOEA.

82. I. *Bonanox*, foliis cordatis acutis integerrimis, caule aculeato, floribus ternis, corollis indiuiſis. M. *S. V.* p. 204. n. 10.
Tanna.

83. I. *carnea*, foliis cordatis glabris: pedunculis multifloris, corollis marginatis. M. *S. V.* p. 205. n. 13.
Societatis inſulae.

CAMPANVLA.

84. C. *gracilis*, foliis lineari-lanceolatis obſolete ſerratis, caule dichotomo, floribus ſolitariis terminalibus. F.
Noua Zeelandia et Noua Caledonia.

NAVCLEA.

85. N. *orientalis*. M. *S. V.* p. 212.
Societatis inſulae.

PORTLANDIA.

86. P. *tetrandra*, floribus tetrandris, foliis obouatis. F.
P. tetrandra. M. *S. V.* p. 213. n. 1.
Inſula ferox (Savage iſland).

SCAEVOLA.

87. S. *Lobelia*. M. *S. V.* p. 213.
Tongatabu.

CINCHONA.

88. C. *corymbifera*, foliis oblongo-lanceolatis, corymbis axillaribus. FORST. in *act. Upſ. nov.* Vol. III. p. 176. M. *S. V.* p. 214. n. 3.

PSYCHOTRIA.

PSYCHOTRIA.

89. P. *ſpecioſa*, arborea, foliis oblongo-lanceolatis, inuolucro terminali ſubtrifloro. F.
Taheiti.

90. P. *aſiatica*, ſtipulis emarginatis, foliis lanceolato-ouatis. M. *S. V.* p. 214. n. 1.
Ibidem.

91. P. *herbacea*, caule herbaceo repente, foliis cordatis petiolatis. M. *S. V.* p. 214. n. 3.
Ibidem.

COFFEA.

92. C. *ſambucina*, foliis oblongo-lanceolatis acutis: cymis corymboſis terminalibus. F.
Inſulae Amicorum.

93. C. *opulina*, foliis ouato lanceolatis; cymis coarctatis globoſis terminalibus. F.
Noua Caledonia.

94. C. *odorata*, foliis ouatis acutis: cymis corymboſis axillaribus. F.
Tanna et Amicorum inſulae.

95. C. *triflora*, foliis ouato-lanceolatis acuminatis: pedunculis terminalibus ternis vnifloris. F.
Taheiti.

CHIOCOCCA.

96. C. *barbata*, erecta, foliis ouatis, pedunculis axillaribus vnifloris, corollis fauce barbatis. F.
Marchionis, Societatis et Amicorum inſulae.

BAEOBOTRYS.

97. B. *nemoralis*. Forst. *Charact. gen.* 11.
Tanna.

DENTELLA.

DENTELLA.

98. D. *repens*. Forst. *Charact. gen.* 13.
Noua Caledonia.

MORINDA.

99. M. *umbellata*, erecta, foliis lanceolato-ouatis, pedunculis confertis. M. *S. V.* p. 217. n. 1.
Inſulae Societatis.

100. M. *citrifolia*, arborea, pedunculis ſolitariis. M. *S. V.* p. 217. n. 2.
Societatis et Amicorum inſulae.

ERITHALIS.

101. E. *polygama*, foliis obouatis: floribus axillaribus, maſculis cymoſis, hermaphroditis ſolitariis. F.
β. Timonius. Rumph. *Amb.* III. p. 216. t. 140. anguſtifolia.
Societatis inſulae.

MVSSAENDA.

102. M. *frondoſa*, panicula foliis coloratis. M. *S. V.* p. 218. n. 1.
Taheiti. Namoka.

GENIOSTOMA.

103. G. *rupeſtris*. Forst. *Charact. gen.* 12.
Tanna.

NICOTIANA.

104. N. *fruticoſa?* foliis lanceolatis ſubpetiolatis amplexicaulibus, floribus acutis, caule fruteſcente. M. *S. V.* p. 221. n. 2.
Botanices inſula prope nouam Caledoniam.

SOLANUM.

105. S. *repandum*, caule inermi ſubherbaceo flexuoſo laeui, foliis ouatis repandis tomentoſis, pedunculis axillaribus cymoſis. F.
Marchionis et Societatis inſulae.

106. S. *nigrum*, caule inermi herbaceo, foliis ouatis dentato-angulatis, racemis diſtichis nutantibus. M. *S. V.* p. 223. n. 20.
Inſula Paſchatis.

107. S. *auiculare*, caule inermi fruticoſo, foliis ſinuato-pinnatifidis: corymbis terminalibus. F.
Noua Zeelandia.

CORDIA.

108. C. *Sebeſtena*, foliis oblongo-ouatis repandis ſcabris. M. *S. V.* p. 230. n. 3.
Societatis inſulae.

109. C. *aſpera*, foliis ouatis acuminatis aſperis: floribus cymoſis rugoſis. F.
Tongatabu.

110. C. *dichotoma*, foliis oblongo-ouatis vix crenatis: corymbis dichotomis. F.
Noua Caledonia.

CARPODETUS.

111. C. *ſerratus*. FORST. *Charact. gen.* 17.
Noua Zeelandia.

CEANOTHUS.

112. C. *capſularis*, foliis ouato-cordatis, acuminatis ſerratis: capſulis triualuibus dehiſcentibus. F.
Taheiti.

CELASTRVS.

113. C. *crenatus*, inermis, foliis ouatis crenulatis: cymis axillaribus. F.
Marchionis inſulae.

CORYNOCARPVS.

114. C. *laeuigata*. Forst. *Charact. gen.* 16. M. *S. V.* p. 242.
Noua Zeelandia.

ARGOPHYLLVM.

115. A. *nitidum*. Forst. *Charact. gen.* 15. M. *S. V.* p. 242.
Noua Caledonia.

ACHYRANTHES.

116. A. *aſpera*, caule fruticoſo erecto, calycibus reflexis ſpicae adpreſſis. M. *S. V.* p. 246. n. 1.
Paſſim intra tropicos.

GYNOPOGON.

117. G. *ſtellatum*, foliis verticillatis ternis lanceolatis Forst. *Charact. gen.* 18. n. 1.
Societatis et Amicorum inſulae.

118. G. *Alyxia*, foliis verticillatis quinis obouatis. F.
Norfolciae inſula.

119. G. *ſcandens*, foliis oppoſitis ouatis coſtatis. F.
Societatis inſulae.

CERBERA.

120. C. *Manghas*, foliis lanceolatis, neruis transuerſalibus. M. *S. V.* p. 251. n. 2.
Societatis inſulae.

121. C. *paruiflora*, foliis ſtellatis obouatis. F.
Inſulae Amicorum et Savage iſland.

GARDENIA.

122. G. *florida*, inermis, corollis obtuſis, calyce angulato, foliis ouatis acutis. M. *S. V.* p. 251. n. 2.
Societatis et Amicorum inſulae.

ECHITES.

123. E. *coſtata*, pedunculis cymoſis, foliis elliptico-lanceolatis acuminatis. F.
Kametti-valli. RHEED. *hort. malab.* P. IX. p. 23. t. 14.
Inſulae Societatis.

TABERNAEMONTANA.

124. T. *citrifolia*, foliis oppoſitis ouatis, floribus lateralibus glomerato-vmbellatis. M. *S. V.* p. 255. n. 1.
Namoka.

DIGYNIA.

MELODINVS.

125. M. *ſcandens*. FORST. *Charact. gen.* 19. M. *S. V.* p. 256.
Noua Caledonia.

PERIPLOCA.

126. P. *capſularis*, foliis lanceolatis integerrimis oppoſitis: cymis axillaribus diffuſis. F.
Noua Zeelandia.

CYNANCHVM.

127. C. *viminale*, caule volubili perenni aphyllo. M. *S. V.* p. 257. n. 1.
Botanices inſula.

ASCLEPIAS.

ASCLEPIAS.

128. A. *volubilis*, arborea volubilis, foliis oppoſitis cordato-ſubrotundis integerrimis carnoſis: vmbellis ſimpliciſſimis erectis. F.
A. volubilis? M. *S. V.* p. 259. n. 4.
Nanſiera-patſia. RHEED. *hort. mal.* P. IX. p. 21. t. 13.
Numularia lactea maior. RUMPH. *amb.* V. t. 175. f. 1.
Tanna.

CHENOPODIVM.

129. C. *triandrum*, foliis cordato-ſagittatis, ſpicis terminalibus aphyllis interruptis. F.
Noua Zeelandia.

SALSOLA.

130. S. *Kali*, herbacea decumbens, foliis ſubulatis ſpinoſis ſcabris, calycibus marginatis axillaribus. M. *S. V.* p. 263. n. 1.
Botanices inſula.

131. S. *fruticoſa*, fruticoſa erecta, foliis filiformibus obtuſiuſculis. M. *S. V.* p. 263. n. 12.
Noua Zeelandia.

GENTIANA.

132. G. *ſaxoſa*, corollis quinquefidis campanulatis, foliis ſpatulatis. F. M. *S. V.* p. 268. n. 14.
Noua Zeelandia.

133. G. *montana*, corollis quinquefidis campanulatis, foliis cordatis ſeſſilibus. F.
Ibidem.

DICHONDRA.

134. D. *repens*. FORST. *Charact. gen.* 20.
Noua Zeelandia.

HYDROCOTYLE.

135. H. *moſchata*, foliis reniformibus ſeptemlobatis ſerratis villoſis: vmbellis multifloris. F.
Noua Zeelandia.

PEVCEDANVM.

136. P. *geniculatum*, foliis ſubrotundo-reniformibus crenatis. F.
Noua Zeelandia.

COPROSVRA.

137. C. *lucida*, pedunculis compoſitis. M. *S. V.* p. 274. n. 1. FORST. *Charact. gen.* 69. n. 2.
Noua Zeelandia.

138. C. *foetidiſſima*, floribus ſolitariis. M. *S. V.* p. 274. n. 2. FORST. *Charact. gen.* 69. n. 1.
Ibidem.

LASERPITIVM.

139. L. *Aciphylla*, caule vaginato, petiolis foliorum dilatatis, foliis digitatis linearibus elongatis mucronatis. M. *S. V.* p. 282. n. 14.
Aciphylla ſquarroſa. FORST. *Charact. gen.* 68.
Noua Zeelandia.

LIGVSTICVM.

140. L. *Gingidium*, foliis pinnatis, foliolis rhombeo-cordatis obliquis crenatis obtuſis. F.
Gingidium montanum. FORST. *Char. gen.* 21.
Noua Zeelandia.

APIVM.

141. A. *graueolens*, foliis caulinis cuneiformibus. M. *S. V.* p. 292. n. 2.
Noua Zeelandia.

TRIGYNIA.

TRIGYNIA.

RHVS.

142\. R. *atrum*, foliis ſimplicibus, ouato-oblongis: floribus polygamis. F.
Noua Caledonia.

XYLOPHYLLA.

143\. X. *longifolia*, foliis linearibus, ramis tetragonis. M. *S. V.* p. 296. n. 1.

PENTAGYNIA.

COMMERSONIA.

144\. C. *echinata.* Forst. *Charact. gen.* 22. M. *S. V.* p. 300.
Societatis inſulae.

LINVM.

145\. L. *monogynum*, foliis alternis lineari-lanceolatis trineruiis, caule ſuffruticoſo. F.
Noua Zeelandia.

POLYGYNIA.

SCHEFFLERA.

146\. S. *digitata.* Forst. *Charact. gen.* 20.
Noua Zeelandia.

CLASSIS VI.

HEXANDRIA.

MONOGYNIA.

CRINVM.

147. C. *aſiaticum?* foliis carinatis. M. *S. V.* p. 318. n. 2.
Amicorum inſulae.

ANTHERICVM.

148. A. *cirratum*, foliis lanceolatis planiuſculis: ſcapo paniculato, filamentis barbatis bicirratis. F.
Noua Zeelandia.

149. A. *Adenanthera*, foliis lineari-enſiformibus baſi connatis; glandula inter filamentum et antheram. F.
Noua Caledonia.

DRACAENA.

150. D. *indiuiſa*, arborea, foliis enſiformibus acutis, racemo (laterali?) compoſito. FORST. *pl. eſculent.* 33.
Noua Zeelandia.

151. D. *auſtralis*, arborea, foliis enſiformibus acutis, racemo terminali erecto ſupradecompoſito. F.
Ibidem.

152. D. *terminalis*, herbacea caulescens, foliis lanceolatis: racemo terminali decompoſito. FORST. *pl. eſculent.* 32.
D. terminalis. M. *S. V.* p. 334. n. 4.
Inſulae Societatis.

PHORMIVM.

PHORMIVM.

153. P. *tenax*, foliis plurimis, infloreſcentia ramoſa. M. *S. V.* p. 336. n. 1. FORST. *Charact. gen.* 24.
Noua Zeelandia.

IVNCVS.

154. I. *campeſtris*, foliis planis ſubpiloſis, ſpicis ſeſſilibus pedunculatisque. M. *S. V.* p. 341. n. 19. γ
Societatis inſulae.

ACHRAS.

155. A. *diſſecta*, floribus confertis corollis octodecemfidis., foliis obouatis emarginato-retuſis. FORST. *pl. eſc.* 13.
A. diſſecta. M. *S. V.* p. 342. n. 3.
Manyl-Kara. RHEED. *hort. mal.* P. III. p. 53. t. 25.
Amicorum Inſulae.

LORANTHVS.

156. L. *tetrapetalus*, pedunculis vnifloris ſubſolitariis, foliis ouatis obtuſis ſubſeſſilibus. F.
L. tetrapetalus. M. *S. V.* p. 343. n. 1.
Noua Zeelandia.

157. L. *Stelis*, racemis trichotomis, pedunculis trigonis, floribus aequalibus. M. *S. V.* p. 344. n. 8.
Societatis inſulae.

DIGYNIA,

GAHNIA.

158. G. *procera*, paniculis ſpicatis pluribus elongatis, floſculis hexandris. F.

G. procera. FORST. *Charact. gen.* 26. M. *S. V.* p. 345.
Noua Zeelandia.

159. G. *Schoenoides*, culmis flexuoſis, paniculis compoſitis ſpicatis rigidis ſubſolitariis. F.
Taheiti.

CLASSIS

CLASSIS VIII.

OCTANDRIA.

MONOGYNIA.

EPILOBIVM.

160. E. *glabellum*, foliis oblongis dentatis obtuſis: imis oppoſitis, caule teretiuſculo erecto. F.
Noua Zeelandia.

161. E. *rotundifolium*, foliis oppoſitis ſubrotundis denticulatis. F.
Ibidem.

XIMENIA.

162. X. *elliptica*, caule inermi, foliis elliptico-lanceolatis: pedunculis multifloris. F.
Noua Caledonia.

SKINNERA.

163. S. *excorticata.* FORST. *Charact. gen.* 29.
Fuchſia excorticata. M. *S. V.* p. 361. n. 3.
Noua Zeelandia.

DODONAEA.

164. D. *viſcoſa*, foliis oblongis. M. *S. V.* p. 362. n. 1.
In Societatis inſulis dioica.
In Noua Zeelandia hermaphrodita.

LAWSONIA.

165. L. *Acronychia*, inermis, foliis longe petiolatis cuneiformibus. M. *S. V.* p. 362. n. 2.
Acronychia laeuis. FORST. *Charact. gen.* 27.
Noua Caledonia.

MELICOPE.

MELICOPE.

166. M. *ternata*. FORST. *Charact. gen.* 28.
Noua Zeelandia.

VACCINIVM.

167. V. *cereum*, pedunculis ſolitariis vnifloris: corollis quinquangulari-ouatis, foliis ouato-ſubrotundis ſerratis. F.
Andromeda cerea. M. *S. V.* p. 406. n. 3.
Bacca, non capſula.
Taheiti.

DAPHNE.

168. D. *foetida*, glabra, floribus terminalibus congeſtis ſeſſilibus: foliis oppoſitis petiolatis ouato-oblongis. M. *S. V.* p. 372. n. 15.
Societatis inſulae.

169. D. *rotundifolia*, hirſuta, floribus terminalibus congeſtis ſeſſilibus, foliis oppoſitis ellipticis ſubpetiolatis glabris. M. *S. V.* p. 372. n. 16.
Namoka.

PASSERINA.

170. P. *Gnidia*, diandra glaberrima, foliis lanceolatis acutis. M. *S. V.* p. 374. n. 1.
Bankſia Gnidia. FORST. *Charact. gen.* 4. n. 1.
Noua Zeelandia.

171. P. *piloſa*, diandra piloſa, foliis linearibus obtuſis. M. *S. V.* p. 374. n. 12.
Bankſia tomentoſa. FORST. *Charact. gen.* 4. n. 2.
Ibidem.

172. P. *proſtrata*, diandra piloſa, foliis ouatis (carnoſis F.). M. *S. V.* p. 374. n. 13.
Bankſia proſtrata. FORST. *Charact. gen.* 4. n. 3.
Ibidem.

DIGYNIA.

DIGYNIA.

WEINMANNIA.

173. W. *racemoſa*, foliis ſimplicibus ouatis ſerratis petiolatis: racemis axillaribus ſolitariis. F.
W. racemoſa. M. *S. V.* p. 376. n. 4.
Noua Zeelandia.

174. W. *paruiflora*, foliis ſimplicibus ouatis ſerratis ſubſeſſilibus: racemis terminalibus ternis quinisue. F.
Taheiti.

CODIA.

175. C. *montana*. Forst. *Charact. gen.* 30. M. *S. V.* p. 376.
Noua Caledonia.

TRIGYNIA.

COCCOLOBA.

176. C. *auſtralis*, foliis cordato-ouatis acutis, floribus polygamis. F.
Noua Zeelandia.

CARDIOSPERMVM.

177. C. *Halicacabum*, foliis laeuibus. M. *S. V.* p. 380. n. 1.
Societatis inſulae.

SAPINDVS.

178. S. *Saponaria*, foliis impari-pinnatis, caule inermi. M. *S. V.* p. 380. n. 1.
Inſula Paſchatis.

TETRAGYNIA.

TETRAGYNIA.

HALORAGIS.

179. H. *proſtrata*, foliis oblongis integerrimis mucronatis, fructibus globoſis. F.
H. proſtrata. M. *S. V.* p. 381. FORST. *Charact. gen.* 31.
Botanices inſula.

180. H. *alata*, foliis ouato-lanceolatis ſerratis, fructibus pyramidatis tetragonis alatis. F. JACQ. *Miſc. Auſtr.* 2. p. 332. *Icon. rarior.* tab. 26.
Cercodia erecta. MVRR. in *Comm. Gotting.* 1780. P. 3. t. 1.
Tetragonia iuaefolia. M. *S. V.* p. 467. n. 3.
Noua Zeelandia.

CLASSIS

CLASSIS IX.

ENNEANDRIA.

MONOGYNIA.

CASSYTA.

181. C. *filiformis*, filiformis laxa. MVRR. *S.V.* p. 384. n. 1.
Societatis et Amicorum inſulae.

CLASSIS X.

DECANDRIA.

MONOGYNIA.

SOPHORA.

182. S. *tomentoſa*, foliis pinnatis, foliolis numeroſis ſubrotundis tomentoſis. M. *S. V.* p. 391. n. 2.
Societatis inſulae.

183. S. *tetraptera*, foliis pinnatis: foliolis numeroſis obouatis emarginatis, leguminibus tetragonis, angulis membranaceis. M. *S. V.* p. 391. n. 7.
Noua Zeelandia.

CASSIA.

184. C. *Sophera*, foliis decemiugis lanceolatis, glandula baſeos oblonga. M. *S. V.* p. 394. n. 28.
Tongatabu.

GVILANDINA.

185. G. *Bonduccella*, aculeata, pinnis oblongo-ouatis, foliolis aculeis geminis. M. *S. V.* p. 396. n. 2.
Noua Zeelandia.

GVAIACVM.

186. G. *dubium*, foliis coniugatis oblongo-lanceolatis obtuſis. F.
Tongatabu.

ADENANTHERA.

ADENANTHERA.

187. A. *ſcandens*, foliis pinnatis biiugis, foliolis ouatis obliquis glabris, cirris terminalibus bifidis. F.
Mallicollo.

TRICHILIA.

188. T. *ſpectabilis*, foliis pinnatis, foliolis obouatis, racemis axillaribus compoſitis. F.
Noua Zeelandia.

189. T. *alliacea*, foliis pinnatis, foliolis lanceolatis acutis: racemis axillaribus ſupradecompoſitis. F.
Namoka.

LIMONIA.

190. L. *minuta*, inermis, foliis pinnatis, racemis corymboſis terminalibus. F.
Amicorum Inſulae.

191. L. *lucida*, inermis, foliis ſimplicibus, pedunculis axillaribus. F.
Mallicollo.

DAIS.

192. D. *diſperma*, floribus octandris decandrisque, foliis ouato-lanceolatis, acuminatis, eneruiis. F.
Tongatabu.

MELASTOMA.

193. M. *Malabathrica*, foliis integerrimis quinqueneruiis lanceolatis ouatis ſcabris. M. *S. V.* p. 404 n. 10.
Societatis inſulae.

194. M. *glabra*, foliis integerrimis trineruiis elliptico-lanceolatis ſcabris, calycibus circumſciſſis. F.
Ibidem.

ANDROMEDA.

195. A. *rupeſtris*, racemis ſimpliciſſimis bracteatis, corollis campanulatis, foliis alternis oblongis ſerrulatis. F.
A. rupeſtris. M. *S. V.* p. 407. n. 16.
Noua Zeelandia.

GAVLTHIERIA.

196. G. *antipoda*, foliis ſparſis ſubrotundis ſerrato-dentatis, caule fruticoſo diffuſo. F.
Noua Zeelandia.

INOCARPVS.

197. I. *edulis*. M. *S. V.* p. 408. FORST. *Charact. gen.* 33.
Societatis, Amicorum et nouarum Hebridum Inſulae.

PENTAGYNIA.

SPONDIAS.

198. S. *dulcis*, petiolis teretibus ſexiugis: foliolis ſerratis coſtatis. F.
Societatis inſulae.

SVRIANA.

199. S. *maritima*. M. *S. V.* p. 431.
Botanices inſula.

CLASSIS

CLASSIS XI.

DODECANDRIA.

MONOGYNIA.

BASSIA.

200. B. *obouata*, foliis obouatis: pedunculis congeſtis terminalibus vnifloris. F.
Tanna.

RHIZOPHORA.

201. R. *gymnorhiza*, foliis ouato-lanceolatis integerrimis: radice terrae ſuperimpoſita. M. *S. V.* p. 442. n. 2.
Namoka.

202. R. *Mangle*, foliis acutis, fructibus ſubulato-clavatis. M. *S. V.* p. 442. n. 5.
Societatis, Amicorum, nouarum Hebridum inſulae et noua Caledonia.

CRATAEVA.

203. C. *religioſa*, inermis, foliolis ſubaequalibus. Forst. *pl. eſc.* 14.
Societatis inſulae.

TRIVMFETTA.

204. T. *procumbens*, foliis ſubrotundo-cordatis ſubtrilobis tomentoſis, caule procumbente. F.
Societatis inſulae.

LYTHRVM.

205. L. *Pemphis*, fruticoſum hirſutum, foliis oppoſitis oblongis integris, floribus axillaribus pedunculatis ſolitariis, capſula circumſciſſa vniloculari. M. *S. V.* p. 447. n. 11.
Pemphis acidula. FORST. *Charact. gen.* 34.
Teautea.

TRIGYNIA.

EVPHORBIA.

206. E. *origanoides*, dichotoma, foliis ſerrulatis ouatis obtuſis trineruiis, panicula terminali, caulibus ſimplicibus. M. *S. V.* p. 449. n. 15.
Amicorum inſulae.

207. E. *Atoto*, dichotoma, foliis ouatis integerrimis, vmbella terminali. F.
Societatis inſulae.

208. E. *glauca*, vmbella ſubquinquefida, trifida, inuolucris et inuolucellis ouatis, foliis ſparſis oblongo-lanceolatis integerrimis, caule fruteſcente. F.
Noua Zeelandia.

TACCA.

209. T. *pinnatifida*. FORST. *Charact. gen.* 35. *pl. eſculent.* 28. M. *S. V.* p. 455.
Societatis Inſulae.

CLASSIS

CLASSIS XII.

ICOSANDRIA.

MONOGYNIA.

MELALEVCA.

210\. M. *ſcoparia*, foliis alternis ouatis acutis ſubtrineruiis: floribus terminalibus ſolitariis, ſeſſilibus. F. *pl. eſc.* 53.
M. ſcoparia. M. *S. V.* p. 699. n. 4.
Leptoſpermum ſcoparium. FORST. *Char. gen.* 36. n. 6.
Noua Zeelandia.

211\. M. *virgata*, foliis oppoſitis linearibus eneruiis, pedunculis axillaribus vmbelliferis. F.
M. virgata. M. *S. V.* p. 699. n. 5.
Leptoſpermum virgatum. FORST. *Charact. gen.* 36. n. 7.
Noua Caledonia.

212\. M. *perforata*, foliis oppoſitis ouali-ſubrotundis, pellucido punctatis, pedunculis axillaribus trifidis. F.
Leptoſpermum perforatum. FORST. *Charact. gen.* 36. n. 4.
Noua Zeelandia.

213\. M. *diffuſa*, foliis oppoſitis ouatis acuminatis: pedunculis rameis cymoſis. F.
Ibidem.

214\. M. *florida*, foliis oppoſitis oblongo-obouatis, venoſis, pedunculis terminalibus cymoſis, calycibus turbinatis. F.

 Leptoſper-

Leptoſpermum ſcandens. FORST. *Charact. gen.* 36. n. 1.
Ibidem.

215. M. *aeſtuoſa*, foliis oppoſitis ouatis obouatisque venoſis (ſubtrineruiis), peduncnlis cymoſis terminalibus tomentoſis. F.
Leptoſpermum collinum. FORST. *Charact. gen.* 36. n. 2.
Melaleuca lucida? M. *S. V.* p. 699. n. 2.
Societatis inſulae.

216. M. *lucida*, foliis oppoſitis lanceolatis vtrinque acuminatis aueniis eneruiis, floribus terminalibus congeſtis ſubſeſſilibus. F.
Noua Zeelandia.

217. M. *ciliata*, foliis ſparſis, ſuperioribus ſuboppoſitis oblongis obtuſis ciliatis aueniis eneruiis: cymis corymboſis terminalibus. F.
M. villoſa? M. *S. V.* p. 699. n. 3.
Leptoſpermum ciliatum. FORST. *Charact. gen.* 36. n. 3.
Noua Caledonia.

218. M. *Leucadendron*, polyadelpha, foliis alternis lanceolatis ſubfalcatis quinqueneruiis, ſpicis terminalibus elongatis. F.
M. Leucadendron. M. *S. V.* p. 698. n. 1. (α latifolia?)

Nomen *Melaleucae*, licet huic vltimae ſpeciei vnice congruum, adoptaui tamen, ne nomina multiplicarentur, ſed genus in locum naturalem reſtitui, cum, praeter M. Leucadendron, nulla ſpecies polyadelpha ſit.
Noua Caledonia.

PSIDIVM.

PSIDIVM.

219. P. *decaſpermum*, foliis ouatis acuminatis pubeſcentibus; pedunculis vnifloris bracteatis. M. *S. V.* p. 461. n. 3.
Decaſpermum fruticoſum. FORST. *Charact. gen.* 37.
Societatis inſulae.

EVGENIA.

220. E. *malaccenſis*, foliis integerrimis, pedunculis ramoſis lateralibus. M. *S. V.* p. 461. n. 1. F. *pl. eſc.* 6.
Paſſim intra tropicos culta.

221. E. *racemoſa*, foliis crenatis, racemis longiſſimis, pomis ouatis quadranguliis. M. *S. V.* p. 461. n. 7.
Noua Caledonia.

TRIGYNIA.

SESVVIVM.

222. S. *Portulacaſtrum.* M. *S. V.* p. 465. n. 1.
Tongatabu.

PENTAGYNIA.

TETRAGONIA.

223. T. *halimifolia*, herbacea papuloſa, foliis elliptico-rhombeis petiolatis, pedunculis axillaribus vnifloris ſubſolitariis, fructu cornuto. F.
T. expanſa? M. *S. V.* p. 467. n. 6.
Noua Zeelandia et Amicorum inſulae.

POLYGYNIA.

RVBVS.

224. R. *auſtralis*, fruticoſus dioicus, foliis ternatis quinato pinnatisque, caule petiolisque aculeatis, racemis axillaribus ſimplicibus. F. Noua Zeelandia.

CLASSIS

CLASSIS XIII.

POLYANDRIA.

MONOGYNIA.

CALOPHYLLVM.

225. C. *Inophyllum*, foliis oualibus. M. *S. V.* p. 491. n. 1.
Societatis inſulae et noua Caledonia.

DICERA.

226. D. *dentata*, monogyna, foliis oblongis acuminatis dentatis, racemis ſimplicibus axillaribus laxis. F.
D. dentata. FORST. *Charact. gen.* 40. n. 1.
Noua Zeelandia.

227. D. *ſerrata*, tetragyna, foliis oppoſitis cordato-ouatis inaequaliter ſerratis: racemis lateralibus compoſitis. F.
D. ſerrata. FORST. *Charact. gen.* 40. n. 2.
Elaeocarpus Dicera. M. *S. V.* p. 494. n. 2.
Ibidem.

TRIGYNIA.

EVRYANDRA.

228. E. *ſcandens.* FORST. *Charact. gen.* 41.
Noua Caledonia.

POLYGYNIA.

WINTERA.

229. W. *axillaris*, pedunculis axillaribus congeſtis vnifloris, piſtillis quatuor. F.
Drimys axillaris. FORST. *Charact. gen.* 42. n. 2. *Nov. Act. Upſ.* III. p. 182. LINN. *Suppl. pl.* p. 270.
Cur LINNEVS junior hanc monogynam fecit, non video.
Noua Zeelandia.

CLEMATIS.

230. C. *hexapetala*, foliis ternatis, foliolis cordato-ouatis crenato-inciſis, pedunculis ramoſis trichotomis diphyllis, floribus polypetalis dioicis. F.
C. hexapetala. M. *S. V.* p. 512. n. 6.
Noua Zeelandia.

231. C. *integrifolia*, foliis ternatis, foliolis ouatis integris mucronatis: pedunculis axillaribus paniculatis diphyllis. F.
Ibidem.

CLASSIS XIV.

DIDYNAMIA.

GYMNOSPERMIA.

TEVCRIVM.

232. T. *villoſum*, foliis elliptico - ouatis acuminatis ſerratis petiolatis villoſis, racemo ſpicato terminali, calycibus inflatis. F.
Tongatabu.

OCYMVM.

233. O. *Scutellarioides*, corollis falcatis, pedicellis ramoſis. M. *S. V.* p. 546. n. 15.
Tanna.

ANGIOSPERMIA.

EVPHRASIA.

234. E. *cuneata*, foliis ſubcuneiformibus inciſis. F.
Noua Zeelandia.

ANTIRRHINVM.

235. A. *hexandrum*, foliis oppoſitis cordato - ouatis ſerratis: pedunculis axillaribus vnifloris. F.
Taheiti.

BESLERIA.

236. B. *biflora*, pedunculis bifloris, inuolucro caduco inflato, foliis ouatis integerrimis. F.
Cyrtandra biflora. FORST. *Charact. gen.* 3. n. 1.
Taheiti.

237.

237. B. *cymoſa*, pedunculis cymoſis, pedicellis bracteolatis, foliis ouatis crenatis. F.
Cyrtandra cymoſa. FORST. *Charact. gen.* 3. n. 2.
Tanna.

MYOPORVM. *Solandr.*

Cal. 5. partitus. Cor. campanulata limbo patente ſubaequali 5 partito. Bacca 1 ſ. 2 ſperma, ſeminibus 2. locularibus.

238. M. *laetum*, foliis oblongis apice ſubſerratis laevibus glabris nitidis, corollis hirſutis. F.
Citharexylum perforatum. Mſpt.
Noua Zeelandia.

239. M. *pubeſcens*, foliis oblongo-ellipticis ſerratis pubeſcentibus. F.
Ibidem.

240. M. *craſſifolium*, foliis oblongis ſubſerratis carnoſis. F.
Botanices inſula.

241. M. *tenuifolium*, foliis lanceolatis acuminatis integerrimis, corollis laeuibus. F.
Citharexylum tenuifolium. Mſpt.
Noua Caledonia.

RVELLIA.

242. R. *reptans*, foliis petiolatis ouatis obtuſis, obtuſe ſerratis, pedunculis terminalibus ſubſpicatis. F.
Tanna.

243. R. *fragrans*, foliis ſeſſilibus oblongis obtuſe ſerratis: floribus axillaribus ſolitariis ſeſſilibus. F.
Taheiti.

VOLKAMERIA.

VOLKAMERIA.

244. V. *inermis*, ramis inermibus. M. *S. V.* p. 577. n. 2.
Paſſim intra tropicos.

VITEX.

245. V. *trifolia*, foliis ternatis quinatisque: foliolis ouatis acutis integerrimis ſubtus caneſcentibus, panicula rachi recta: pedicellis dichotomis. M. *S. V.* p. 579. n. 5.
Amicorum inſulae.

AVICENNIA.

246. A. *reſiniſera*, foliis ouato-lanceolatis ſubtus tomentoſis. FORST. *pl. eſculent.* 44.
Noua Zeelandia.

ACANTHVS.

247. A. *ilicifolius*, foliis repandis dentato-ſpinoſis, caule fruticoſo aculeato. M. *S. V.* p. 580. n. 5.
Tanna, Noua Caledonia.

CLASSIS

CLASSIS XV.

TETRADYNAMIA.

SILICVLOSA.

LEPIDIVM.

248. L. *oleraceum*, foliis elliptico-oblongis acutis ſerratis: floribus tetrandris. FORST. *pl. eſc.* 38.
Noua Zeelandia.

249. L. *Piſcidium*, foliis elliptico-oblongis acutis integerrimis: floribus tetradynamis. FORST. *pl. eſculent.* 39.
Botanices inſula, Teautea, Huaheine.

SILIQVOSA.

SISYMBRIVM.

250. S. *heterophyllum*, foliis pinnatis, foliolis reniformibus ſubtrilobis, infimis pinnatifidis, piloſis. F.
Noua Zeelandia.

CLASSIS

CLASSIS XVI.

MONADELPHIA.

PENTANDRIA.

WALTHERIA.

251. W. *indica*, foliis ouatis ſerratis plicatis: capitulis (ſub-)ſeſſilibus. M. *S. V.* p. 610. n. 3.

252. W. *Lophanthus*, foliis ſubrotundo-cordatis ſerratis ſericeo-tomentoſis petiolatis: capitulis pedunculatis imbricato-bractatis. F.
Lophanthus tomentoſus. FORST. *Charact. gen.* 14.
Marchionis inſulae.

MELOCHIA.

253. M. *odorata*, cymis corymboſis axillaribus, foliis cordatis acuminatis ſerratis. F.
M. odorata. M. *S. V.* p. 611. n. 7.
Tanna, Ea-uwa.

DECANDRIA.

PLAGIANTHVS.

254. P. *diuaricatus*. FORST. *Charact. gen.* 43.
Noua Zeelandia.

POLYANDRIA.

BARRINGTONIA.

255. B. *ſpecioſa*. FORST. *char. gen.* 38. M. *S. V.* p. 680.
Societatis et Amicorum inſulae.

SIDA

SIDA.

256. S. *rhombifolia*, foliis lanceolato-rhomboidalibus ſerratis, axillis ſubſpinoſis. M. *S.V*. p. 621. n. 4.
Amicorum nouarumque Hebridum inſulae.

257. S. *indica*, foliis cordatis ſublobatis, ſtipulis reflexis, pedunculis petiolo longioribus, capſulis multilocularibus ſcabris, calyce longioribus. M. *S V*. p. 623. n. 24.
Tanna. Noua Caledonia.

URENA.

258. U. *lobata*, foliis angulatis. M. *S.V*. p. 627. n. 1.
Paſſim intra tropicos.

GOSSYPIVM.

259. G. *religioſum*, foliis trilobis acutis ſubtus vniglanduloſis, ramulis nigropunctatis. M. *S.V*. p. 628. n. 4.
Societatis inſulae.

HIBISCVS.

260. H. *bacciferus*, foliis cordatis integerrimis acuminatiſſimis, calyce exteriore triphyllo caduco, fructu baccato. F.
H. populneus? M. *S.V*. p. 629. n. 10.
Societatis et Amicorum inſulae, etiam Paſchatis inſula.

261. H. *tiliaceus*, foliis cordatis ſubrotundis indiuiſis acuminatis, crenatis, caule arboreo, calyce exteriore decemdentato. M. *S.V*. p. 629. n. 11. FORST. *pl. eſc*. 45.
Paſſim intra tropicos.

262. H. *Roſaſinenſis*, foliis ouatis acuminatis ſerratis, caule arboreo. M. *S.V*. p. 629. n. 13.
Ibidem.

263. H. *Abelmoſchus*, foliis ſubpeltato-cordatis ſeptemangularibus, caule hiſpido. M. *S. V.* p. 630. n. 26.
Societatis inſulae.

264. H. *eſculentus?* foliis quinquepartito-pedatis, calycibus interioribus latere rumpentibus. M. *S. V.* p. 630. n. 27.
Ibidem.

265. H. *haſtatus*, foliis oblongis ſubhaſtato-trilobis integerrimis, pedunculis axillaribus terminalibusque ſubtrifidis: petalis laceris. F.
H. haſtatus. M. *S. V.* p. 631. n. 33.
Ibidem.

CROSSOSTYLIS.

266. C. *biflora*. FORST. *Char. gen.* 44.
Inſulae Societatis.

CLASSIS XVII.

DIADELPHIA.

DECANDRIA.

ABRVS.

267. A. *precatorius.* M. *S.V.* p. 641. n. 1.
Societatis inſulae.

ERYTHRINA.

268. E. *Corallodendron*, foliis ternatis inermibus, caule arboreo inermi. M. *S.V.* p. 642. n. 2. *β.*
Societatis inſulae.

DOLICHOS.

269. D. *luteolus*, volubilis, leguminibus capitatis pluribus cylindricis, ſeminibus rotundatis. M. *S.V.* p. 657. n. 5.
Societatis inſulae.

270. D. *pruriens*, volubilis, leguminibus racemoſis, valuulis ſubcarinatis hirtis, pedunculis ternis? M. *S.V.* p. 657. n. 12.
Namoka.

GLYCINE.

271. G. *roſea*, volubilis, foliis ternatis, foliolis cordato-ouatis acutis integerrimis, pedunculis trifloris: leguminibus compreſſis carinatis monoſpermis. F.
Inſulae Societatis.

272.

272. G. *lucida*, volubilis, foliis ternatis, foliolis ouatis acuminatis integerrimis, racemis terminalibus glanduloſis, pedicellis ternis, leguminibus inflatis diſpermis. F.
Ibidem.

AESCHYNOMENE.

273. A. *coccinea*, caule arboreo, foliis pinnatis, foliolis ouali-oblongis obtuſis mucronatis, leguminibus ſubarcuato-filiformibus compreſſiusculis. F.
A. coccinea. M. *S. V.* p. 671. n. 3.
Toeri-Mera. RUMPH. *amb.* I. t. 77.
Differentia ſpecifica A. grandiflorae emendanda, vt ab hac diſtinguatur.
Societatis inſulae, Botanicesque inſula.

HEDYSARVM.

274. H. *vmbellatum*, foliis ternatis, pedunculis vmbelliferis, caule fruticoſo. M. *S. V.* p. 673. n. 22.
Tanna et noua Caledonia.

275. H. *heterocarpon*, foliis ternatis, floribus paniculato-ſpicatis, leguminibus articulatis, infimo monoſpermo, ſtipulis ſetaceis. M. *S. V.* p. 673. n. 24.
Societatis inſulae.

276. H. *Lagopodioides*, foliis ternatis, racemis oblongis, leguminibus inflexis, calycibus hirſutis. M. *S. V.* p. 674. n. 40.
Noua Caledonia.

GALEGA.

277. G. *littoralis*, leguminibus racemoſis, tota villoſo-tomentoſa. M. *S.V.* p. 679. n. 3.
Paſſim intra tropicos.

LOTVS.

278. L. *arboreus*, leguminibus quinatis, foliolis obcordatis, caule arboreo. F.
Noua Zeelandia.

CLASSIS XVIII.

POLYADELPHIA.

ICOSANDRIA.

CITRVS.

279. C. ***Aurantium***, petiolis alatis, foliis acuminatis. M. *S. V.* p. 679. n. 2.
Tanna.

280. C. ***decumana***, petiolis alatis, foliis obtuſis emarginatis. M. *S. V.* p. 679. n. 3.
Amicorum Inſulae.

POLYANDRIA.

HYPERICVM.

281. H. ***gramineum***, floribus trigynis, caule herbaceo tetragono dichotomo, foliis oblongo-parabolicis.
Noua Caledonia.

CLASSIS XIX.

SYNGENESIA.

POLYGAMIA AEQVALIS.

SONCHVS.

282. S. *oleraceus*, pedunculis tomentoſis, calycibus glabris. M. *S.V.* p. 712. n. 5.
Noua Zeelandia et Amicorum inſulae.

BIDENS.

283. B. *piloſa*, foliis pinnatis ſubpiloſis, caule geniculis barbatis, calycibus inuolucro ſimplici, ſeminibus diuergentibus. M. *S.V.* p. 732. n. 7.
Tongatabu.

ADENOSTEMMA.

284. A. *viſcoſa*. FORST. *Charact. gen.* 45.
Verbeſina Lavenia. M. *S.V.* p. 779. n. 4.
Societatis inſulae.

AGERATVM.

285. A. *conyzoides*, foliis ouatis, caule piloſo. M. *S.V.* p. 737. n. 1.
Tanna.

CHRYSOCOMA.

286. C. *purpurea*, herbacea, foliis elliptico-lanceolatis ſubſerratis pubeſcentibus, panicula terminali corymboſa. F.
Tanna.

CALEA.

CALEA.

287. C. *leptophylla*, floribus terminalibus ternis quinisue congeſtis, foliis oblongis imbricatis integerrimis ſeſſilibus, caule fruticoſo. F.
Noua Zeelandia.

288. C. *pinifolia*, pedunculis terminalibus congeſtis, foliis linearibus aceroſis, caule fruticoſo. F.
Ibidem.

POLYGAMIA SVPERFLVA.

GNAPHALIVM.

289. G. *trineruе*, fruticoſum, foliis ſeſſilibus lanceolatis trineruiis ſubtus argenteis, panicula corymboſa terminali laxa. F.
Noua Zeelandia.

290. G. *lanatum*, herbaceum, lanatum, foliis lineari-oblongis apice calloſo-mucronatis, corymbo coarctato, caule ſimpliciſſimo. F.
Ibidem.

291. G. *inuolucratum*, herbaceum, foliis linearibus elongatis mucronatis ſubtus tomentoſis, capitulo terminali folioſo. F.
Ibidem.

292. G. *luteo-album*, herbaceum, foliis ſemiamplexicaulibus enſiformibus repandis obtuſis vtrinque pubeſcentibus. M. *S. V.* p. 747. n. 32.
Noua Caledonia.

XERANTHEMVM.

293. X. *Bellidioides*, (receptaculo nudo, pappo piloſo) ramis vnifloris, foliis ouato-oblongis cuſpidatis amplexicaulibus, pedunculis tomentoſis ſubaphyllis. F.
Noua Zeelandia.

CINERARIA.

294. C. *rotundifolia*, paniculis paucifloris, foliis petiolatis ouato-ſubrotundis integerrimis ſubtus tomentoſis, caule arboreo. F.
Brachyglottis rotundifolia. FORST. *Charact. gen.* 46. n. 1.
Noua Zeelandia.

295. C. *repanda*, paniculis compoſitis racemoſis diffuſis, foliis petiolatis ouatis repando-ſinuatis, ſubtus tomentoſis, caule arboreo. F.
Brachyglottis repanda. FORST. *Charact. gen.* 46. n. 2.
Ibidem.

ASTER.

296. A. *holoſericeus*, herbaceus, foliis oblongo-lanceolatis ſerratis ſubtus argenteo-ſericeis, ſcapis vnifloris folioſis. F.
Noua Zeelandia.

297. A. *coriaceus*, herbaceus, foliis ouatis integerrimis ſuperne ſulcatis ſubtus lanatis, ſcapis vnifloris folioſis lanatis. F.
Ibidem.

SOLIDAGO.

298. S. *arboreſcens*, caule arboreo, paniculis corymboſis ſtrictis terminalibus, foliis ouato-ſubrotundis acutis repandis petiolatis. F.
Noua Zeelandia.

ARNICA.

299. A. *oporina*, fruticoſa, foliis lanceolatis calloſo-crenatis ſubtus tomentoſis, pedunculis vnifloris ſolitariis terminalibus ſquamatis. F.
Noua Zeelandia.

COTVLA.

COTVLA.

300. C. *coronopifolia*, foliis lanceolato-linearibus amplexicaulibus dentatis, floribus floſculoſis. M. *S.V.* p. 774. n. 4.
Noua Zeelandia.

301. C. *minuta*, foliis alternis ſubamplexicaulibus oblongis dentatis, floribus ſolitariis ſeſſilibus folio oppoſitis. F.
Noua Caledonia.

ECCLIPTA.

302. E. *proſtrata*, caule proſtrato, foliis ſubundulatis ſubpetiolatis. M. *S.V.* p. 779. n. 4.
Tanna.

SIGESBECKIA.

303. S. *orientalis*, petiolis ſeſſilibus, calycibus exterioribus linearibus maioribus patentibus. M. *S.V.* p. 779. n. 1.
Paſſim intra tropicos.

BVPHTHALMVM.

304. B. *Helianthoides*, foliis oppoſitis ouatis ſerratis triplineruiis, calycibus folioſis, caule herbaceo. M. *S.V.* p. 781. n. 11.
Paſſim intra tropicos.

POLYGAMIA NECESSARIA.

CALENDVLA.

305. C. *pumila*, ſeminibus oblongis incuruatis, foliis orbiculatis crenato-ſerratis, ſerraturis mucronatis, ſcapo nudo vnifloro. F.
Noua Zeelandia.

POLYGAMIA SEGREGATA.

CRASPEDIA.

Calyculus nullus. Calyx communis imbricatus. Floſculorum faſciculi aliquot depreſſi, omnes hermaphroditi tubuloſi. Pappus plumoſus. Receptaculum paleaceum.

306. C. *vniflora.* F.
Noua Zeelandia.

MONOGAMIA.

SHAWIA.

307. S. *paniculata.* FORST. *Charact. gen.* 48.
Noua Zeelandia.

LOBELIA.

308. L. *arborea*, caule arboreo, foliis petiolatis ouato-oblongis ſerrato-dentatis, pedunculis axillaribus ſolitariis vnifloris. F.
Societatis Inſulae.

309. L. *angulata*, caule proſtrato ſimpliciſſimo, foliis ſubrotundis repando-dentatis, pedunculis axillaribus ſolitariis vnifloris. F.
Noua Zeelandia.

CLASSIS

CLASSIS XX.

GYNANDRIA.

DIANDRIA.

OPHRYS.

310. O? *ſquamata*, bulbis faſciculatis, ſcapo elongato aphyllo ſpicato, foliis radicalibus imbricatis oblongis acutis carinatis, nectarii labio trifido barbato deflexo. F.
Noua Caledonia.

311. O. *vnifolia*, bulbo ouato, ſcapo tereti vaginato, folio tereti fiſtuloſo reflexo, in medio pro emittendo ſcapo perforato. F.
Noua Zeelandia.

SERAPIAS.

312. S. *regularis*, bulbis ouatis fibroſis, foliis vaginantibus enſiformibus carinatis, ſcapo erecto ſpicato, corollis hexapetalis. F.
Thelymitra longifolia. FORST. *Charact. gen.* 49.
Suaſu b. SOLANDRI huc reduxi.
Noua Zeelandia.

EPIDENDRVM.

313. E. *tuberoſum?* foliis lato-lanceolatis neruoſis membranaceis, bulbo innatis, ſcapo vaginato, nectario cymbiformi bifido. M. *S. V.* p. 819. n. 25.
Noua Caledonia.

314. E. *triſte*, foliis teretibus cylindricis inanibus, vaginis fiſtuloſis, pedunculis oppoſitifoliis corymboſis vaginam perforantibus, nectarii labio integro ſpatulato cordiformi. F.
Ibidem.

315. E. *criſpatum*, foliis caulinis teretibus filiformibus ſubincuruis inanibus ſeſſilibus: racemo ſimpliciſſimo, nectarii labio lacinia media reuoluta, vndulato crenata. F.
Societatis inſulae.

316. E. *equitans*, foliis equitantibus enſiformi-compreſſis carinatis, acuminatis, ſcapo nudo, racemo ſpicato nutante filiformi. F.
Ibidem.

317. E. *Myoſurus*, foliis baſi vaginantibus oblique divaricatis linearibus obtuſis canaliculatis, ſcapo nudo, racemo ſpicato nutante filiformi. F.
Ibidem.

318. E. *biflorum*, foliis diſtichis, lineari lanceolatis acutis, pedunculis bifloris ſolitariis vaginas foliorum perforantibus, nectarii labio trifido acuminato. F.
Ibidem.

319. E. *autumnale*, foliis caulinis lanceolatis neruoſis confertis vaginantibus, pedunculis terminalibus ſquamatis paniculatis, nectarii labio erecto retuſo. F.
Noua Zeelandia.

320. E. *Faſciola*, foliis radicantibus paraſiticis faſciculatis linearibus vndulatis, ſcapis pluribus racemoſis, nectarii labio calceiformi calcarato. F.
Societatis inſulae.

321. E. *vmbellatum*, ſtolonibus repentibus, foliis oualibus carnoſis petiolatis, petiolis infixis apophyſibus

phyſibus ouatis anguloſo-carinatis, ſcapo radicato, vmbella ſimplici vnilaterali, petalis infimis elongatis. F.
Ibidem.

322. E. *reſupinatum*, foliis caulinis petiolatis ouatis acuminatis integerrimis plicato carinatis quinqueneruiis, racemo ſimplici ſpicato, floribus reſupinatis, nectarii labio inferiore dentato. F. Baſaala-poulou-maravara. RHEED. *hort. mal.* Part. XII. tab. 27. bona.
Societatis inſulae.

323. E. *Clypeolum*, folio radicali ſubrotundo-cordato, baſi cucullato, apice reflexo acuto, ſcapo erectiuſculo racemoſo, nectarii labio inferiore orbiculato emarginato maximo. F.
Ibidem.

FORSTERA.

324. F. *ſedifolia.* M. *S. V.* p. 819. n. 1. *Nov. Act. Upſ.* III. p. 184. t. 9.
Noua Zeelandia.

TRIANDRIA.

SISYRINCHIVM.

325. S. *Ixioides*, foliis enſiformibus conduplicatis longiſſimis, panicula trichotoma, petalis exterioribus dimidio minoribus. F.
Noua Zeelandia.

PENTANDRIA.

PENTANDRIA.

PASSIFLORA.

326. P. *aurantia*, foliis trilobis, lobis parabolicis diſtantibus, medio productiore. F.
Noua Caledonia.

POLYANDRIA.

GREWIA.

327. G. *Mallococca*, foliis cordatis ouato-oblongis crenatis ſcabris, pedicellis axillaribus trifloris, fructu tetracocco. M. *S. V.* p. 827. n. 4.
Malococca crenata. FORST. *Charact. gen.* 39. *Nov. Act. Upſ.* III. p. 185.
Societatis et Amicorum inſulae.

ARVM.

328. A. *eſculentum*, acaule, foliis peltatis ouatis integerrimis baſi emarginatis. M. *S. V.* p. 827. n. 7. FORST. *pl. eſculent.* 26.
Paſſim intra tropicos et in Noua Zeelandia colitur.

329. A. *macrorhizon*, acaule, foliis peltatis cordatis repandis baſi bipartitis. M. *S. V.* p. 827. n. 8. FORST. *pl. eſc.* 27.
Arum indicum ſatiuum. RUMPH. *amb.* V. t. 106.
Ibidem.

DRACONTIVM.

330. D. *polyphyllum*, ſcapo breuiſſimo, petiolo radicato lacero, foliolis tripartitis, laciniis pinnatifidis.

natifidis. M. *S. V.* p. 829. n. 1. Forst. *pl. eſculent.* 29.

Societatis inſulae planta culta.

331. D. *pertuſum*, foliis pertuſis, caule ſcandente. M. *S. V.* p. 829. n. 5.

Tanna.

CLASSIS

CLASSIS XXI.

MONOECIA.

MONANDRIA.

ARTOCARPVS.

332. A. *inciſa*, foliis inciſis. M. *S. V.* p. 838. n. 1. Monogr. mea *vom Brodbaum* 1784. 4. Forst. *pl. eſc.* 1.
Paſſim intra tropicos culta.

BALANOPHORA.

333. B. *fungoſa.* Forst. *Charact. gen.* 50.
Tanna.

CASVARINA.

334. C. *equiſetifolia*, verticillis ſtaminum approximatis. M. *S. V.* p. 840. n. 1.
Intra tropicos.

335. C. *nodiflora*, verticillis ſtaminum remotis. M. *S. V.* p. 840. n. 2.
Noua Caledonia.

TRIANDRIA.

TYPHA.

336. T. *latifolia*, foliis enſiformibus, ſpica maſcula femineaque approximatis. M. *S. V.* p. 841. n. 1.
Noua Zeelandia.

COIX.

COIX.

337. C. *Lacryma.* M. *S. V.* p. 842.
Tongatabu.

CAREX.

338. C. *vncinata*, ſpica ſimplici androgyna lineari ſuperne maſcula, ariſtis feminarum vncinatis, maſculis muticis. M. *S. V.* p. 843. n. 5.
Noua Zeelandia.

339. C. *lithoſperma*, ſpicis paniculatis maſculis, floribus femineis ſolitariis, ſeminibus globoſis nitidis. M. *S. V.* p. 846. n. 45.
Tanna.

HERNANDIA.

340. H. *ſonora*, foliis peltatis. M. *S. V.* p. 847. n. 1.
Societatis et Amicorum inſulae.

PHYLLANTHUS.

341. P. *virgata*, foliis ſimplicibus alternis linearibus mucronatis, pedunculis axillaribus ſolitariis vnifloris, caule fruticoſo. F.
Societatis inſulae.

TETRANDRIA.

URTICA.

342. U. *aeſtuans*, foliis alternis cordatis, racemis dichotomis, fructibus orbiculato-corymboſis. M. *S. V.* p. 850. n. 16.
Societatis inſulae.

343. U. *argentea*, foliis alternis elliptico-lanceolatis integris ſubtus glaucis: ſpicis axillaribus ſolitariis interruptis. F.
Ibidem.

344. U. *ruderalis*, foliis alternis cordato-ouatis obtuſe ſerratis, paniculis axillaribus diuaricato-corymboſis. F.
Ibidem.

345. U. *virgata*, foliis oppoſitis ouatis ſerratis trinerviis, ſpicis axillaribus ſolitariis interruptis. F.
Ibidem.

346. U. *ferox*, foliis oppoſitis haſtato-cordatis ſetoſo-ſerratis, racemis geminis diuaricatis, ſtipulis cordatis. F.
Noua Zeelandia.

MORVS.

347. M. *papyrifera*, foliis palmatis, fructibus hiſpidis M. *S. V.* p. 851. n. 3.
Culta intra tropicos paſſim, etiam in Noua Zeelandia borealiore.

PENTANDRIA.

AMARANTHVS.

348. A. *tricolor*, glomerulis triandris axillaribus ſubrotundis ſeſſilibus, foliis lanceolatis acuminatis. M. *S. V.* p. 853. n. 4.
Societatis inſulae.

349. A. *gangeticus*, glomerulis triandris ſubſpicatis ouatis, foliis laoceolato-ouatis emarginatis. M. *S. V.* p. 853. n. 6.
Ibidem.

HEPTANDRIA.

GVETTARDA.

350. G. *ſpecioſa*. M. *S. V.* p. 855. n. 1.
Societatis inſulae.

MONADEL-

MONADELPHIA.

CVPRESSVS.

351. C. *columnaris*, foliis imbricatis ſubulatis ſulcatis, ſtrobilis cylindricis elongatis. F.
Noua Caledonia et Norfolciae inſula.

ACALYPHA.

352. A. *virgata*, ſpicis femineis inuolucris cordatis ſerratis, maſculis diſtinctis aphyllis, foliis lanceolatis ouatis. M. *S. V.* p. 862. n. 2.
Tanna.

CROTON.

353. C. *variegatum*, foliis lanceolatis integerrimis glabris pictis petiolatis. M. *S. V.* p. 863. n. 1.
Mallicollo et Tanna.

354. C. *nutans*, foliis rhombeo-ſubrotundis acuminatis vndulatis glabris nutantibus. F.
Societatis, Amicorum et nouarum Hebridum Inſulae.

355. C. *Inophyllum*, foliis obouatis integerrimis, caule arboreo. F.
Noua Caledonia.

RICINVS.

356. R. *Mappa*, foliis peltatis indiuiſis. M. *S. V.* p. 866. n. 4.
Tanna.

357. R. *dioicus*, foliis cordatis acuminatis. F.
Ibidem.

STERCVLIA.

358. S. *Balanghas*, foliis ouatis integerrimis petiolatis floribus paniculatis. M. *S. V.* p. 866. n. 1. Forst. *pl. eſculent.* 22.
Tanna.

359. S. *foetida*, foliis digitatis. M. *S. V.* p. 866. n. 2. Forst. *pl. eſc.* 23.
Ibidem.

ALEVRITES.

360. A. *triloba.* Forst. *Charact. gen.* 56.
Inſulae Societatis.

SYNGENESIA.

GLOCHIDION.

361. G. *ramiflorum.* Forst. *Charact. gen.* 57.
Societatis inſulae et nouae Hebrides.

CVCVRBITA.

362. C. *lagenaria*, foliis ſubangulatis tomentoſis, baſi ſubtus biglanduloſis, pomis lignoſis. M. *S. V.* p. 868. n. 1.
Intra tropicos et in Paſchatis inſula.

SICYOS.

363. S. *angulata*, foliis angulatis. M. *S. V.* p. 870. n. 1.
Noua Zeelandia.

CLASSIS XXII.

DIOECIA.

MONANDRIA.

ASCARINA.

364. A. *polyſtachya*. FORST. *Charact. gen.* 59.
Societatis inſulae.

PANDANUS.

365. P. *odoratiſſima*. M. *S. V.* p. 878.
Athrodactylis ſpinoſa. FORST. *Charact. gen.* 75.
Paſſim intra tropicos.

TRIANDRIA.

MABA.

366. M. *elliptica*. FORST. *Charact. gen.* 61. M. *S. V.* p. 881.
Amicorum inſulae.

RESTIO.

367. R. *ſimplex*, culmis ſimplicibus, ſpica terminali. M. *S. V.* p. 882. n. 6.
Noua Zeelandia.

EXCOECARIA.

368. E. *Agallocha*. M. *S. V.* p. 882.
Tongatabu.

TETRANDRIA.

VISCVM.

369. V. *opuntioides*, caule prolifero ramoſiſſimo aphyllo compreſſo. M. *S. V.* p. 883. n. 4.
Societatis inſulae.

370. V. *antarcticum*, foliis ouatis integerrimis, racemis rameis terminalibusque articulatis. F.
Noua Zeelandia.

PENTANDRIA.

MELICYTVS.

371. M. *ramiflorus*. Forst. *Charact. gen.* 62.
Noua Zeelandia.

HEXANDRIA.

SMILAX.

372. S. *Ripogonum*, caule inermi tereti radicante, foliis ouato-lanceolatis acuminatis quinqueneruiis, floribus hermaphroditis. F.
Ripogonum ſcandens. Forst. *Charact. gen.* 25.
Noua Zeelandia.

373. S. *purpurata*, caule inermi tereti dichotomo, foliis cordato-ouatis acuminatis vnguiculatis integerrimis quinqueneruiis, pedunculis axillaribus vmbelliferis. F.
Noua Caledonia.

DIOSCOREA.

374. D. *pentaphylla*, foliis digitatis. M. *S. V.* p. 888. n. 1.
Societatis inſulae.

375. D. *alata*, foliis cordatis, caule alato bulbifero. M. *S.V.* p. 888. n. 5.
Paſſim culta.

376. D. *bulbifera*, foliis cordatis, caule laeui bulbifero. M. *S.V.* p. 888. n. 6.
Societatis et Marchionis inſulae.

DECANDRIA.

CORIARIA.

377. C. *ſarmentoſa*, procumbens diffuſa, foliis cordato-ouatis acuminatis integerrimis quinqueneruiis ſubpetiolatis: racemis axillaribus elongatis nutantibus. F.
Noua Zeelandia.

DODECANDRIA.

MENISPERMVM.

378. M. *japonicum*, foliis peltatis rotundato-ouatis integris. M. *S.V.* p. 892. n. 3. THVNB. *flor. japon.* p. 193.
Inſulae Societatis.

POLYANDRIA.

HEDYCARYA.

379. H. *dentata.* FORST. *Char. gen.* M. *S.V.* p. 894.
Noua Zeelandia.

XYLOSMA.

380. X. *ſuaueolens*, foliis ouatis ſerratis. F.
Myroxylon ſuaueolens. Forst. *Charact. gen.* 63. n. 1.
Nomen mutaui, ne confundatur cum Myroxylo M. *S. V.* p. 395. quod Balſamum peruuianum fundit.
Inſulae Societatis.

381. X. *orbiculatum*, foliis ſubrotundis integerrimis. F.
Myroxylon orbiculatum. Forst. *Charact. gen.* 63. n. 2.
Savage iſland.

CLASSIS

CLASSIS XXIII.

POLYGAMIA.

MONOECIA.

MVSA.

382. M. *paradiſiaca*, ſpadice nutante. M. *S. V.* p. 902. n. 1.
Paſſim intra tropicos culta.

ISCHAEMVM.

383. I. *muticum*, ſeminibus muticis. M. *S. V.* p. 906. n. 1.
Tanna.

384. I. *murinum*, ſpica bipartita: calyce ſeminibusque ariſtatis. Forst. *Nov. Act. Upſ.* III. p. 185.
Ibidem.

385. I. *inuolutum*, ſpica ſecunda mutica quadriflora, inuoluta receptaculo foliaceo concauo. F.
Societatis inſulae et paſſim intra tropicos.

CENCHRVS.

386. C. *echinatus*, ſpica oblonga conglomerata. M. *S. V.* p. 907. n. 5.
Societatis inſulae.

PARIETARIA.

387. P. *debilis*, foliis alternis ouatis petiolatis integerrimis ſubpiloſis, pedunculis axillaribus ſubtrifloris, caule erectiuſculo. F.
Noua Zeelandia.

TERMINALIA.

388. T. *Catappa*, foliis obouatis, ſubtus tomentoſis. M. *S.V.* p. 910. n. 1.
Tanna.

389. T. *glabrata*, foliis obouatis vtrinque glabris. Forst. *pl. eſc.* 20.
Societatis et Amicorum inſulae.

CLVSIA.

390. C. *pedicellata*, foliis oppoſitis obouatis integerrimis venoſis, cymis axillaribus, floribus tetrapetalis. F.
Noua Caledonia.

391. C. *ſeſſilis*, foliis oppoſitis obouatis ellipticisque integerrimis venoſis, floribus axillaribus ſolitariis ſubſeſſilibus tetrapetalis. F.
Tongatabu.

POMETIA.

392. P. *pinnata*, foliis pinnatis, racemo ſupradecompoſito terminali. F.
P. pinnata. Forst. *Charact. gen.* 55.
Tanna, Namoka.

393. P. *ternata*, foliis ternatis, racemis ſimpliciuſculis axillaribus. F.
Aporetica ternata. Forst. *Charact. gen.* 66.
Noua Caledonia.

CELTIS.

394. C. *orientalis*, foliis oblique cordatis ſubtus villoſis. M. *S.V.* p. 912. n. 3.
Societatis Inſulae.

MIMOSA.

MIMOSA.

395. M. ***Mangium***, inermis arborea, foliis ouatis integerrimis neruoſis obtuſis: ſpicis globoſis pedunculatis. L.
M. ſimplicifolia. M. *S. V.* p. 913. n. 1.
Mangium montanum. RUMPH. *amb.* III. t. 81.
Inſulae Amicorum, nouae Hebrides, noua Caledonia.

DIOECIA.

PENNANTIA.

396. P. *corymboſa.* FORST. *Charact. gen.* 67.
Noua Zeelandia.

PISONIA.

397. P. ***inermis***, caule inermi. M. *S. V.* p. 920. n. 2.
Societatis inſulae.

PANAX.

398. P. ***arboreum***, foliis quinatis obouatis ſerrato-dentatis, vmbellis compoſitis. F.
P. arborea. M. *S. V.* p. 920. n. 4.
Noua Zeelandia.

399. P. *ſimplex*, foliis alternis lanceolatis ſerratis, vmbellis compoſitis. F.
Ibidem.

BREYNIA.

400. B. *diſticha.* FORST. *Charact. gen.* 73.
Noua Caledonia, Tanna.

GRISELINIA.

401. G. *lucida.* F.
Scopolia lucida. FORST. *Charact. gen.* 70.
Noua Zeelandia.

TRIOECIA.

TRIOECIA.

FICVS.

402. F. *religioſa*, foliis cordato ouatis integerrimis acuminatiſſimis, transuerſe ſtriatis: fructibus geminis globoſis ſeſſilibus calyculatis. F. F. religioſa. M. *S. V.* p. 921. n. 4.
Tanna.

403. F. *ſcabra*, foliis cordato-ouatis obliquis integris ſubtus ſcabris: fructibus turbinatis ecalyculatis. F.
Tanna, Namoka.

404. F. *aſpera*, foliis oblique cordatis ſinuato-dentatis vtrinque aſperis: fructibus turbinatis, calycis margine obſoleto adnato. Forst. *pl. eſculent.* 7.
Tanna.

405. F. *tinctoria*, foliis oblique ouatis obtuſis, fructibus turbinatis baſi calyculatis. F.
Societatis inſulae.

406. F. *indica?* foliis lanceolatis integerrimis petiolatis, pedunculis aggregatis, ramis radicantibus. M. *S. V.* p. 922. n. 7. Forst. *pl. eſc.* 9.
Inſulae Amicorum, nouarum Hebridum, noua Caledonia.

407. F. *ſeptica*, foliis obliquis oblongo-ouatis acuminatis: pedunculis geminis apice calyculatis, fructibus verrucoſis. F.
Tanna.

408. F. *Granatum*, foliis ouatis integerrimis, pedunculis terminalibus geminis horizontaliter divergentibus, fructibus calyculatis globoſis. Forst. *pl. eſculent.* 8.
Ibidem.

409.

409. F. *obliqua*, foliis lanceolatis glaberrimis margine cartilagineis: pedunculis geminis breuiſſimis, calycibus caducis longitudine fructus. F.
Namoka, Tanna.

410. F. *prolixa*, foliis lanceolato-oblongis acuminatis ſubtus punctatis: pedunculis geminis apice calyculatis. F.
Societatis inſulae.

CLASSIS

CLASSIS XXIV.

CRYPTOGAMIA.

FILICES.

CYCAS.

411. C. *circinalis*, frondibus pinnatis, foliolis linearibus planis. M. *S. V.* p. 925. n. 1.
Amicorum et nouarum Hebridum Inſulae.

OPHIOGLOSSVM.

412. O. *ſcandens*, caule flexuoſo tereti, frondibus coniugatis pinnatis, foliolis vtrinque ſpiciferis. M. *S. V.* p. 926. n. 7.
Paſſim intra tropicos.

OSMVNDA.

413. O. *diſcolor* ****, frondibus pinnatis, pinnis oblongis acutiuſculis integris ſeſſilibus alternis approximatis. F.
Noua Zeelandia.

414. O. *procera* ****, frondibus pinnatis, pinnis remotis ouato-oblongis acuminatis ſerratis ſeſſilibus. F.
Ibidem.

ACROSTICHVM.

415. A. *dichotomum*, nudum dichotomum, ſpicis ſecundis adſcendentibus reflexis. M. *S. V.* p. 929. n. 12.
Societatis inſulae.

416.

416. A. *aureum*, fronde pinnata: pinnis alternis linguiformibus integerrimis glabris. M. *S. V.* p. 929. n. 16.
Ibidem.

PTERIS.

417. P. *pedata*, frondibus quinquangulis trifoliatis, pinnis pinnatifidis: lateralibus bipartitis. M. *S. V.* p. 931. n. 15.
Inſulae Societatis.

418. P. *eſculenta*, frondibus ſupradecompoſitis ſulcatis, foliolis pinnatis, pinnis linearibus decurrentibus, ſummis breuioribus. FORST. *pl. eſculent.* 47.
Ibidem.

419. P. *comans*, frondibus pinnatis: foliolis pinnatifidis, pinnis elongatis lanceolatis, apice attenuatis ſerratis. F.
Noua Zeelandia.

420. P. *rotundifolia*, frondibus pinnatis hiſpidis: pinnis ſuboppoſitis ſubrotundis, obſolete crenatis. F.
Ibidem.

421. P. *humilis*, frondibus ſubbipinnatis: foliolis oblongis inciſis ſubpinnatis, extimis obſolete crenatis confluentibus. F.

BLECHNVM.

422. B. *orientale*, frondibus pinnatis: pinnis linearibus alternis. M. *S. V.* p. 931. n. 2.
Societatis inſulae.

HEMIONITIS.

423. H. *reticulata*, frondibus lanceolato-falcatis integerrimis, venis reticulatis. F.
Societatis inſulae.

LONCHITIS.

LONCHITIS.

424. L. *tenuifolia*, arboreſcens, frondibus decompoſitis, foliis pinnatis: pinnis lineari oblongis ſerratis, infimis pinnatifidis. F.
Tanna.

ASPLENIVM.

425. A. *Nidus*; frondibus ſimplicibus lanceolatis integerrimis glabris. M. *S. V.* p. 933. n. 4.
Societatis inſulae.

426. A. *flaccidum*, frondibus pinnatis: foliolis alternis remotis pinnatifidis, ſegmentis linearibus ſtrictis. F.
Noua Zeelandia.

427. A. *lucidum*, frondibus pinnatis: foliolis oppoſitis oblongo-ouatis acuminatis ſerrulatis. F.
Ibidem.

428. A. *polyodon*, frondibus pinnatis: foliolis trapezoidibus acuminatis acutis duplicato-ſerratis. F.

429. A. *obliquum*, frondibus pinnatis, ſtipitibus ſquamuloſis: foliolis oblongis oppoſitis acuminatis ſerratis, margine exteriore breuiore. F.
Noua Zeelandia.

430. A. *obtuſatum*, frondibus pinnatis: foliolis oppoſitis oblongis obtuſis ſerratis. F.
Ibidem.

431. A. *tenerum*, frondibus pinnatis: foliolis rhombeo-oblongis, obtuſis inciſo-ſerratis. F.

432. A. *caudatum*, frondibus pinnatis: foliolis pinnatifidis linearibus apice ſetaceis, ſegmentis obtuſis apice inciſo-ſerratis, ſtipite hirto. F.

433. A. *bulbiferum*, frondibus ſubbipinnatis, foliolis decurrentibus oblongis obtuſis pinnatifidis, fructificationibus proliferis. F.
Noua Zeelandia.

POLYPODIVM.

434. P. *Acroſtichoides*, frondibus linearibus integris glabris, fructificationibus confertis. F.
Societatis inſulae.

435. P. *ſerpens*, frondibus linearibus obtuſis pubeſcentibus, fructificationibus ſerialibus, ſtolonibus repentibus. F.
Noua Zeelandia.

436. P. *puſtulatum*, frondibus pinnatifidis laeuibus, pinnis oblongis integris acuminatis. F.

437. P. *ſcandens*, frondibus pinnatifidis laeuibus: pinnis linearibus obtuſis vndulatis diſtantibus, ſtolonibus radicantibus ſcandentibus. F.
Societatis inſulae.

438. P. *euectum*, frondibus pinnatis, pinnis oppoſitis oblongis apice lineari-acuminatis ſerratis, fructificationum ſerie ſubmarginali continua. F.
Ibidem.

439. P. *hirſutulum*, frondibus pinnatis: pinnis oblongis obtuſe ſerratis, baſi ſurſum auriculatis, ſtipite villoſo. F.

440. P. *tenellum*, frondibus pinnatis: foliolis alternis remotis linearibus acuminatis vndulatis. F.

441. P. *diſſectum*, frondibus pinnatis: foliolis pinnatifidis, pinnulis oblongis obtuſe ſerratis, inferioribus longioribus vix coalitis. F.

442. P. *nymphale*, frondibus pinnatis pubeſcentibus: foliolis linearibus acuminatiſſimis ſerratis, baſi pinnatifidis: ſegmentis oblongis obtuſis vix antrorſum falcatis. F.
Noua Zeelandia.

443. P. *inuiſum*, frondibus pinnatis glabris: foliolis linearibus acuminatiſſimis ſerrato-pinnatis:

pinnis lanceolatis falcato-acutis, baſi connatis. F.

Ibidem.

444. P. *pennigerum*, frondibus pinnatis glabris: foliolis linearibus acuminatiſſimis ſubpinnatis: pinnis ouato-oblongis obtuſis connatis. F.

Ibidem.

445. P. *veſtitum*, frondibus ſubbipinnatis: pinnis rhombeo-ouatis acutis inciſo-ſerratis, infimis lobatis ſubpinnatis, ſtipite ſquamis ſcarioſis veſtito. F.

446. P. *nudum*, frondibus bipinnatis: foliolis pinnatis: pinnis rhombeis inciſo-crenatis, ſtipite ſcabro nudo. F.

447. P. *ſetoſum*, frondibus bipinnatis: foliolis ſubpinnatis: pinnis linearibus inciſo-ſerratis, ſerraturis ſetaceis, ſtipite villoſo. F.

448. P. *ariſtatum*, frondibus bipinnatis: foliolis inferioribus pinnatis: pinnis rhombeo-oblongis inciſis, ſegmentis mucronato-ſerratis, ſtipite villoſiuſculo. F.

449. P. *adiantiforme*, frondibus ſubbipinnatis: foliolis ouatis inciſis, lobis ouatis obtuſis crenato-ſerratis, infimis diſcretis, ſtipite ſquamuloſo aſpero. F.

450. P. *dichotomum*, frondibus dichotomis bipinnatis: pinnulis linearibus integris parallelis. M. *S. V.* p. 938. n. 66. FORST. *pl. eſc.* 49.

Societatis inſulae et Noua Zeelandia.

451. P. *arboreum?* frondibus bipinnatis ſerratis, caudice arboreo inermi. M. *S. V.* p. 938. n. 67.

Ibidem.

452. P. *medullare*, frondibus bipinnatis: foliolis pinnatis acuminatiſſimis: pinnis oblongis ſubfalcatis

falcatis acutis crenatis, ſtipite aſpero, caudice arboreo hiſpido. FORST. *pl. eſc.* 48.
Noua Zeelandia.

453. P. *extenſum*, frondibus bipinnatis: foliolis pinnatis acuminatis apice ſerratis: pinnis oblongis ſerratis, ſtipite punctis aſpero, caudice arboreo. F.

454. P. *dealbatum*, frondibus bipinnatis: foliolis pinnatis acuminatis ſubtus albis: pinnis oblongis ſubfalcatis ſerratis: ſtipite aſpero, caudice arboreo. F.
Noua Zeelandia.

455. P. *affine*, frondibus bipinnatis: foliolis pinnatis acuminatis: pinnis lineari-oblongis, crenatis, ſtipite glabro. F.

456. P. *lunulatum*, frondibus bipinnatis: foliolis pinnatis apice ſerratis ſetaceis, pinnis lineari-oblongis falcatis apice ſerrulatis, ſtipite aſpero. F.

457. P. *latifolium*, frondibus ſubbipinnatis: foliolis ouatis acuminatis pinnatifidis lobatisque: laciniis repandis crenatis, ſtipite glaberrimo nitido. F.

ADIANTVM.

458. A. *pedatum*, frondibus pedatis: foliolis pinnatis: pinnis antice gibbis inciſis fructificantibus. M. *S. V.* p. 939. n. 5.
Societatis inſulae.

459. A. *clauatum*, frondibus ſupradecompoſitis: foliolis alternis: pinnis cuneiformibus integerrimis alternis vnifloris. M. *S. V.* p. 940. n. 22.

460. A. *trapeziforme*, frondibus ſupradecompoſitis: foliolis alternis: pinnis rhombeis inciſis vtrinque fructificantibus. M. *S V.* p.940. n. 24?
Noua Zeelandia et intra tropicos.

461. A. *cuneatum*, frondibus pinnatis: foliolis oppoſitis ſubpinnatis: pinnis cuneiformibus retuſis alternis. F.

TRICHOMANES.

462. T. *reniforme*, frondibus ſimplicibus reniformibus ſtipitatis multifloris, receptaculis ſeminum exſertis cylindricis. F.
Noua Zeelandia.

463. T. *contiguum*, frondibus pinnatis: pinnis linearibus alternis approximatis integris, fructificantibus apice inciſis. F.

464. T. *humile*, frondibus pinnatifidis dichotomis: pinnis alternis decurrentibus linearibus obtuſis integris, fructificationibus turbinato-infundibuliformibus, ſtylis ſetaceis exſertis, ſtipite vix vllo. F.
Societatis inſulae.

465. T. *ſanguinolentum*, frondibus ſubbipinnatis, foliolis alternis pinnatifidis: pinnis dichotomis lineari-oblongis obtuſis integris decurrentibus: fructificationibus ouato-ſubrotundis dehiſcentibus. F.
Noua Zeelandia.

466. T. *biualue*, frondibus ſubbipinnatis: pinnis alternis decurrentibus dichotomis, ſegmentis linearibus ſerratis, fructificationibus ſubrotundis biualuibus. F.

467. T. *dilatatum*, frondibus ſubbipinnatis: pinnis alternis dichotomis decurrentibus cuneiformibus inciſis, fructificationibus biualuibus orbicularibus inflatis. F.

468. T. *demiſſum*, frondibus bipinnatis: foliolis alternis ſtrictis pinnatis: pinnis pinnatifido-dichotomis, ſegmentis linearibus obtuſis integris: fructificationibus globoſis terminalibus. F.

469. T. *cuneiforme*, frondibus bipinnatis: foliolis alternis dichotomis pinnatis: pinnis alternis cuneiformibus inciſis, fructificationibus truncato-laceris. F.

470. T. *gibberoſum*, frondibus bipinnatis: foliolis pinnatis: pinnis oblongis pinnatifido-inciſis, ſegmentis linearibus integris, margine interiore infra apicem fructificantibus. F.

471. T. *epiphyllum*, frondibus bipinnatis: foliolis pinnatis acuminatis: pinnis linearibus inciſo-ſerratis: fructificationibus in pagina ſuperiore infra apicem inciſurarum ſquamaceis. F.

472. T. *flaccidum*, frondibus decompoſitis: foliolis rhombeis ſubpinnatis: pinnis cuneiformibus apice inciſis margine interiore infra apicem fructificantibus. F.

473. T. *multifidum*, frondibus decompoſitis: foliolis alternis pinnatis: pinnis dichotomis linearibus decurrentibus argute ſerratis: fructificationibus ouato-ſubrotundis dehiſcentibus. F.

474. T. *elatum*, frondibus decompoſitis: foliolis pinnatis: pinnis oblongis ſtrictis pinnatifido-inciſis, ſegmentis oblongis apice ſerratis, fructificationibus terminalibus ouatis. F.

475. T. *ſolidum*, frondibus decompoſitis: foliolis acuminatis: pinnis ouato-oblongis inciſis crenatis: fructificationibus filiformibus tubuloſis. F.

476. T. *ſquarroſum*, frondibus ſubbipinnatis: foliolis oblongis acuminatis ſubpinnatis ſerratis, ſerraturis mucronatis: fructificationibus ſinuum globoſis dehiſcentibus ſubbiualuibus, ſtipite hiſpido barbato, caudice arboreo. F.

MVSCI.

LYCOPODIVM.

477. L. *nudum*, foliis ſubnullis, ſpicis dichotomis: floribus diſtantibus. M. *S. V.* p. 942. n. 2.
Societatis inſulae.

478. L. *Phlegmaria*, foliis verticillato-quaternis, ſpicis terminalibus dichotomis. M. *S. V.* p. 942. n. 2.
Ibidem.

479. L. *ſquarroſum*, foliis ſparſis linearibus glabris, ſpicis terminalibus folioſis. F.

480. L. *cernuum*, foliis ſparſis curuatis, caule ramoſiſſimo, ſpicis nutantibus. M. *S. V.* p. 943. n. 12.

481. L. *complanatum*, foliis bifariis connatis, ſuperficialibus ſolitariis, ſpicis geminis pedunculatis. M. *S. V.* p. 943. n. 18.

482. L. *volubile*, foliis bifariis diſtichis falcatis acuminatis, caule volubili ramoſo hiſpido, pedunculis ramoſis, ſpicis aggregatis nudis. F.

483.

483. L. *flabellatum*, foliis bifariis: ſuperficialibus diſtichis, caule erecto tereti. M. *S. V.* p. 943. n. 23.

484. L. *ſcarioſum*, foliis bifariis imbricatis erectis ouatis glabris, ſpicis terminalibus ſquamoſis, ſquamis ſcarioſis reflexis. F.

485. L. *myrtifolium*, foliis quadrifariis lineari-oblongis, ſpicis terminalibus filiformibus folioſis. F.

APPENDIX.

PALMAE.

CORYPHA.

486. C. *umbraculifera*, frondibus pinnato-palmatis plicatis filoque interiectis, ſtipitibus ciliato-ſpinoſis. M. *S. V.* p. 984. n. 1.
Marchionis et Amicorum inſulae.

COCOS.

487. C. *nucifera*, inermis, frondibus pinnatis: foliolis replicatis enſiformibus. M. *S. V.* p. 985. n. 1.
Paſſim intra tropicos.

ARECA.

488. A. *oleracea??* foliolis integerrimis. M. *S. V.* p. 986. n. 2.
Nouae Hebrides; forte ad A. ſapidam referenda.

PLANTAE OBSCVRAE. †)

I.

489. *Salicornia auſtralis.* S. Noua Zeelandia.

 II.

†) Hoc titulo colliguntur plantae in Auſtraſiae inſulis a nobis quidem lectae, ſed vel imperfecte, vel omnino, deficientibus floribus et fructificationibus, non deſcriptae. Nomina nonnullis deinde adiecta iuxta ſententiam b. Solandri, qui easdem olim deſcripſerat.

II.

490. *Iaſminum.* Noua Caledonia.

III.

491. *Schoenus arundinaceus.* S. Noua Caledonia.
492. *Schoenus ſurinamenſis.* S. Taheiti.
493. *Schoenus rubiginoſus.* S. Noua Zeelandia.
494. *Schoenus eleuatus.* S. Societatis inſulae.
495. *Cyperus venuſtus.* S. Ibidem.
496. *Cyperus ſtuppeus.* S. Ibidem.
497. *Cyperus linctus.* S. Ibidom.
498. *Poa caeſpitoſa.* Noua Zeelandia.
499. *Poa imbecilla.* S. Ibidem.

IV.

500. *Galium vmbroſum.* S. Noua Zeelandia.
501. *Nertera depreſſa.* S. Ibidem.

V.

502. *Menyanthes.* Hort. Malab. Part. XI. t. 29. Noua Caledonia.
503. *Conuoluulus chryſorrhizus.* S. Forst. *pl. eſculent.* 24. Paſſim cultus.
504. *Scaeuola ſericea.* Botanices inſula ad nouam Caledoniam.
505. *Scaeuola ſaligna.* Noua Caledonia.
506. *Phyſalis flaccida.* S. Societatis Inſulae.
507. *Solanum viride.* S. Forst. *pl. eſculent.* 42. Ibidem.
508. *Solanum aſtroites.* Ibidem.
509. *Rhamnus poliſolius.* S. Noua Zeelandia.

510. *Rhamnus Zizyphoides.* S. Societatis Inſulae.
511. *Rondeletioides.* S. Ibidem.
512. *Hydrocotyle.* Noua Caledonia.
513. *Coproſma puſilla.* Noua Zeelandia.

VI.

514. *Iuncus tenax.* S. Noua Zeelandia.
515. *Rumex flexuoſus.* S. Ibidem.

VIII.

516. *Epilobium junceum.* S. Ibidem.
517. *Polygonum imberbe.* S. Societatis inſulae.
518. *Polyſcias pinnata.* FORST. *Charact. gen.* 32. Tanna.

X.

519. *Oxalis reptans.* S. Societatis inſulae et Noua Zeelandia.

XI.

520. *Portulaca lutea.* S. Societatis inſulae.
521. *Euphorbia.* Norfolciae inſula.

XII.

522. *Eugenia? paniculata.* Tongatabu.
523. *Meſembryanthemum auſtrale.* S. Noua Zeelandia.

XIII.

524. *Ranunculus riuularis.* S. Noua Zeelandia.
525. *Ranunculus hirtus.* S. Ibidem.

XIV.

XIV.

526. *Stachys decemdentata.* S. Societatis inſulae.

527. *Ocymum puſillum.* Tanna.

528. *Scrophularioides arborea.* Societatis et Marchionis inſulae.

XV.

529. *Cardamine ſarmentoſa.* S. Teautea.

XVI.

530. *Geranium patulum.* S. Noua Zeelandia.

531. *Geranium piloſum.* S. Ibidem.

532. *Sida.* Ibidem.

XVII.

533. *Phaſeolus amoenus.* S. Societatis inſulae.

XIX.

534. *Scorzonera ſcapigera.* S. Noua Zeelandia.

535. *Gnaphalium redolens.* S. Noua Caledonia.

536. *Baccharis.* Noua Caledonia.

537. *Baccharis* II. Ibidem.

538. *Senecio lautus.* S. Noua Zeelandia.

539. *Senecio anguſtifolius.* S. Ibidem.

540. *Cotula bicolor.* S. Tanna.

541. *Buphthalmum vniflorum.* Norfolciae inſula.

542. *Buphthalmum procumbens.* Amicorum inſulae.

543. *Coreopſis Helianthoides.* Intra tropicos.

544. *Coreopſis fruticoſa.* Ibidem.

545. *Coreopſis.* Tanna.

XX.

546. *Epidendrum diſtichum.* Societatis inſulae.

547. *Epidendrum.* Ibidem.

XXI.

548. *Carex hamata.* Noua Zeelandia.

549. *Carex ternaria.* S. Ibidem.

550. *Carex debilis.* S. Ibidem.

551. *Croton clutioides.* Tanna.

552. *Croton.* Tanna.

553. *Glochidion.* Tanna.

554. *Cucurbita pruriens.* S. Societatis inſulae.

555. *Cucurbita aſpera.* S. Ibidem.

556. *Cucurbita multiflora.* S. Ibidem.

XXII.

557. *Maba maior.* FORST. *pl. eſculent.* 21. Amicorum inſulae.

558. *Meryta lanceolata.* FORST. *Charact. gen.* 60. Societatis inſulae.

559. *Meriſta laeuigata.* S. Noua Zeelandia.

560. *Cliffortioides cordata.* S. Noua Zeelandia.

361. *Dacrydium cupreſſinum.* S. FORST. *pl. eſcul.* 54. Ibidem.

XXIII.

562. *Andropogon.* Tanna.

563. *Holcus redolens.* S. Noua Zeelandia.

564. *Ixalum inerme.* S. Ibidem.

565. *Mimoſa glanduloſa.* S. Societatis inſulae.

566. *Mimoſa.* Noua Caledonia.

567. *Mimoſa* II. Ibidem.

568. *Mimoſa* III. Ibidem.

569. *Ceodes umbellata.* FORST. *Charact. gen.* 71. Tanna.

570. *Meliſtaurum diſtichum.* FORST. *Charact. gen.* 72. Noua Caledonia.

571. *Ficus ſtipulacea.* Tanna.

XXIV.

572. *Oſmunda reptans.* S. Noua Zeelandia.

573. *Oſmunda obtuſata.* S. Ibidem.

574. *Aſplenium polyanthes.* S. Societatis inſulae.

575. *Bryum.*

576. *Bryum* II.

577. *Bryum* III.

578. *Hypnum.*

579. *Hypnum* II.

580. *Hypnum* III.

581. *Hypnum* IV.

582. *Hypnum* V.

583. *Iungermannia.*

584. *Lichen.*

585. *Lichen* II.

586. *Lichen* III.

587. *Lichen* IV.

588. *Lichen* V.

589. *Fucus.*

590. *Fucus* II.

591. *Ulva.*

APPENDIX.

APPENDIX.

592. *Areca ſapida.* S. Noua Zeelandia. Forst. *pl. eſc.* 35.

593. *Mawhaha* radix. Cook *it.* I. 332. 393. Forst. *pl. eſc.* 31. Amicorum inſulae.

594. *Dgidgi* (Jeejee) radix. Cook l. c. Forst. *pl. eſc.* 30. Ibidem.

INDEX.

INDEX.

A

Abrus precatorius No. 267
Acalypha virgata 352
Acanthus ilicifolius 247
Achras diſſecta 155
Achyranthes aſpera 116
Acroſtichum aureum 416
— — dichotomum 415
Adenanthera ſcandens 187
Adenoſtemma viſcoſa 284
Adianthum clauatum 459
— — cuneatum 461
— — pedatum 458
— — trapeziforme 460
Aeſchynomene coccinea 273
Ageratum Conyzoides 285
Agroſtis ouata 40
Aira antarctica 41
Aleurites triloba 360
Amaranthus gangeticus 349
— — tricolor 348
Amomum Curcuma 2
— — Zerumbet 1
Anciſtrum diandrum 52
Andromeda rupeſtris 195
Andropogon 562
Anthericum Adenanthera 149
— — cirratum 148
Anthoxanthum crinitum 18
Antirrhinum hexandrum 235
Apium graueolens No. 141
Areca oleracea 488
— ſapida 592
Argophyllum nitidum 115
Arnica oporina 299
Artocarpus inciſa 332
Arum eſculentum 328
— macrorhizon 329
Arundo Bambos 47
— conſpicua 48
Aſcarina polyſtachya 364
Aſclepias volubilis 128
Aſplenium bulbiferum 433
— — caudatum 432
— — flaccidum 426
— — lucidum 427
— — nidus 425
— — obliquum 429
— — obtuſatum 430
— — polyanthes 574
— — polyodon 428
— — tenerum 431
Aſter coriaceus 297
— holoſericeus 296
Auena filiformis 46
Auicennia reſinifera 246

B

Baccharis 536
— ij 537

Baeobotrys

Baeobotrys nemoralis 97
Balanophora fungoſa 333
Barringtonia ſpecioſa 255
Baſſia obouata 200
Beſleria biflora 236
— cymoſa 237
Bidens piloſa 283
Blackburnia pinnata 53
Blechnum orientale 422
Boerhaavia erecta 4
— — tetrandra 5
Breynia diſticha 400
Bryum 575
— II 576
— III 577
Buphthalmum Helianthoides 304
— — procumbens 542
— — vniflorum 541

C

Calea leptophylla 287
— pinifolia 288
Calendula pumila 305
Calophyllum Inophyllum 225
Campanula gracilis 84
Cardamine ſarmentoſa 529
Cardioſpermum Halicacabum 177
Carex debilis 550
— hamata 548
— lithoſperma 339
— ternaria 549
— vncinata 338
Carpodetus ſerratus 111
Caſſia Sophera 184
Caſſyta filiformis 181
Caſuarina equiſetifolia 334
— — nodiflora 335

Ceanothus capſularis No. 112
Celaſtrus crenatus 113
Celtis orientalis 394
Cenchus echinatus 386
Ceodes vmbellata 569
Cerbera Manghas 120
— — paruiflora 121
Chenopodium triandrum 129
Chiococca barbata 96
Chryſocoma purpurea 286
Cinchona corymbifera 88
Cineraria repanda 295
— — rotundifolia 294
Citrus Aurantium 279
— decumana 280
Clematis hexapetala 230
— — integrifolia 231
Cliffortioides cordata 560
Cluſia pedicellata 390
— ſeſſilis 391
Coccoloba auſtralis 176
Cocos nucifera 487
Codia montana 175
Coffea odorata 94
— — opulina 93
— — ſambucina 92
— — triflora 95
Coix Lacryma 337
Commelina virginica? 26
Commerſonia echinata 144
Conuoluulus braſilienſis 81
— — chryſorrhizus 503
— — coeleſtis 77
— — corymboſus 80
— — grandiflorus? 76
— — mucronatus 79
— — peltatus 78

Conuoluulus

Conuoluulus tuguriorum No. 74
— — turpethum 75
Coproſma foetidiſſima 138
— — lucida 137
— — puſilla 513
Cordia aſpera 109
— dichotoma 110
— Sebeſtena 108
Coreopſis 545
— fruticoſa 544
— helianthoides 543
Coriaria ſarmentoſa 377
Corynocarpus laeuigata 114
Corypha vmbraculifera 486
Cotula bicolor 540
— coronopifolia 300
— minuta 301
Craſpedia vniflora 306
Crataeua religioſa 203
Crinum aſiaticum 147
Croſſoſtylis biflora 266
Croton 552
— clutioides 551
— ſnophyllum 355
— nutans 354
— variegatum 353
Cucurbita aſpera 555
— — lagenaria 362
— — multiflora 556
— — pruriens 554
Cupreſſus columnaris 351
Cycas circinalis 411
Cynanchum viminale 127
Cynoſurus indicus 45
Cyperus linctus 497
— odoratus 27
— ſtuppeus 396
Cyperus venuſtus No. 495

D

Dais diſperma 192
Daphne foetida 168
— rotundifolia 169
Dentella repens 98
Dianthera caerulea 14
— — clauata 15
Dgidgi (radix) 594
Dicera dentata 226
— ſerrata 227
Dichondra repens 134
Dioſcorea alata 375
— — bulbifera 376
— — pentaphylla 374
Dodonaea viſcoſa 164
Dolichos luteolus 269
— pruriens 270
Dorſtenia lucida 58
— pubeſcens 59
Dracaena auſtralis 151
— — indiuiſa 150
— — terminalis 152
Dracontium pertuſum 331
— — polyphyllum 330

E

Echites coſtata 123
Ecclipta proſtrata 302
Embothrium vmbellatum 60
Epacris faſcicularis 72
— juniperina 71
— longifolia 68
— pumila 70
— roſmarinifolia 69

G Epidendrum.

Epidendrum No. 547
— — autumnale 319
— — biflorum 318
— — clypeolum 323
— — criſpatum 315
— — diſtichum 546
— — equitans 316
— — faſciola 320
— — myoſurus 317
— — reſupinatum 322
— — triſte 314
— — tuberoſum 313
— — vmbellatum 321
Epilobium glabellum 160
— — junceum 516
— — rotundifolium 161
Erithalis obouata 101
Erythrina Corallodendron 268
Eugenia malaccenſis 220
— paniculata 522
— racemoſa 221
Euphorbia 521
— — Aloto 207
— — glauca 208
— — origanoides 206
Euphraſia cuneata 234
Euryandra ſcandens 228
Excoecaria Agallocha 368

F

Fagara Euodia 54
Ficus aſpera 404
— Granatum 408
— indica 406
— obliqua 409
— prolixa 410
— religioſa 402
Ficus ſcabra No. 403
— ſeptica 407
— ſtipulacea 571
— tinctoria 405
Forſtera ſedifolia 324
Fucus 589
— II. 590

G

Gahnia procera 158
— Schoenoides 159
Galega littoralis 277
Galium vmbroſum 500
Gardenia florida 122
Gaulthieria antipoda 196
Genioſtoma rupeſtris 103
Gentiana ſaxoſa 132
— — montana 133
Geranium patulum 530
— — piloſum 531
Glochidion 553
— — ramiflorum 361
Glycine lucida 272
— roſea 271
Gnaphalium inuolucratum 291
— — lanatum 290
— — luteo-album 298
— — redolens 535
— — trinerue 289
Goſſypium religioſum 259
Gratiola Monnieria 16
Grewia Mallococca 327
Griſelinia lucida 401
Guaiacum dubium 186
Guettarda ſpecioſa 350
Guilandina Bonduccella 185
Gynopogon Alyxia 118

Gyriopogon ſcandens No. 119
— — ſtellatum 117

H

Haloragis alata 180
— proſtrata 179
Hedycarya dentata 379
Hedyſarum heterocarpon 275
— — lagopodioides 276
— — vmbellatum 274
Hemionitis reticulata 423
Hernandia ſonora 340
Hibiſcus Abelmoſchus 263
— — bacciferus 260
— — eſculentus 264
— — haſtatus 265
— — Roſa ſinenſis 262
— — tiliaçeus 261
Holcus redolens 563
Hydrocotyle 512
— — moſchata 135
Hypericum gramineum 281
Hypnum 578
— II 579
— III 580
— IV 581
— V 582

I

Iaſminum 490
— — didymum 8
— — ſimplicifolium 7
Inocarpus edulis 197
Ipomoea Bona nox 82
— carnea 83
Iſchaemum inuolutum 385
— — murinum 384
Iſchaemum muticum No. 383
Iuncus campeſtris 154
— tenax 514
Iungermannia 583
Iuſticia longifolia 13
— repanda 12
Ixalum inerme 564

K

Kyllinga monocephala 50
— — triceps 31

L

Laſerpitium Aciphylla 139
Lawſonia Acronychica 165
Lepidium oleraceum 248
— — piſcidium 249
Lichen 584
— II 585
— III 586
— IV 587
— V 588
Liguſticum Gingidium 140
Limonia lucida 191
— minuta 190
Linum monogynum 145
Lithoſpermum incanum 63
Lobelia angulata 307
— arborea 308
Lonchitis tenuifolia 424
Loranthus Stelis 157
— — tetrapetalus 156
Lotus arboreus 278
Lycopodium cernuum 480
— — complanatum 481
— — flabellatum 483
— — myrtifolium 485

Lycopodium nudum No. 477
— — phlegmaria 478
— — ſcarioſum 484
— — ſquarroſum 479
— volubile 482
Lyſimachia decurrens 65
Lythrum Pemphis 205

M

Maba elliptica 366
— maior 557
Mawhaha (radix) 593
Melaleuca aeſtuoſa 215
— — ciliata 217
— — diffuſa 213
— — florida 214
— — Leucadendron 218
— — lucida 216
— — perforata 212
— — ſcoparia 210
— — virgata 211
Melaſtoma glabra 194
– – malabathrica 193
Melicope ternata 166
Melicytus ramiflorus 371
Meliſtaurum diſtichum 570
Melochia odorata 253
Melodinus ſcandens 125
Meniſpermum japonicum 378
Menyanthes 502
Meriſta laeuigata 559
Meryta lanceolata 558
Meſembryanthemum auſtrale 523
Mimoſa 566
— II 567
— III 568
— glandulοſa 565
Mimoſa Mangium 393
Mniarum biflorum 6
Morinda citrifolia 100
— vmbellata 99
Morus papyrifera 347
Muſa paradiſiaca 382
Muſſaenda frondoſa 102
Myoporum craſſifolium 240
— — laetum 238
— — pubeſcens 239
— — tenuifolium 241
Myoſotis ſpatulata 62

N

Nauclea orientalis 85
Nertera depreſſa 501
Nicotiana fruticoſa? 104

O

Ocymum puſillum 527
— — Scutellarioides 233
Oldenlandia debilis 56
— — foetida 55
— — tenuifolia 57
Ophiorrhiza ſubumbellata 66
Ophiogloſſum ſcandens 412
Ophrys ſquamata 310
— vnifolia 311
Oſmunda diſcolor 413
— — obtuſata 573
— — procera 414
— — reptans 572
Oxalis reptans 519

P

Panax arboreum 398
— ſimplex 399

Pandanus odoratiſſima No. 365
Panicum compoſitum 39
— — filiforme 38
— — hiſpidum 36
— — ſanguinale 37
Parietaria debilis 387
Paſpalum orbiculare 35
Paſſerina Gnidia 170
— — piloſa 171
— — proſtrata 172
Paſſiflora aurantia 326
Pennantia corymboſa 396
Periploca capſularis 126
Peteſia carnea 51
Peucedanum geniculatum 136
Phalaris canarienſis 34
Phaſeolus amoenus 533
Phormium tenax 153
Phyllanthus virgata 341
Phyſalis flaccida 506
Piper acuminatum? 23
— excelſum 20
— latifolium 22
— methyſticum 21
— pallidum 24
— Siriboa 19
— tetraphyllum 25
Piſonia inermis 397
Plagianthus diuaricatus 254
Plumbago Zeylanica 73
Poa anceps 43
— caeſpitoſa 498
— Eragroſtis? 42
— imbecilla 499
— latifolia 44
Polygonum imberbe 517
Polypodium Acroſtichoides 434
Polypodium adiantiforme No. 449
— — affine 455
— — arboreum 451
— — ariſtatum 448
— — dealbatum 454
— — dichotomum 450
— — diſſectum 441
— — euectum 438
— — extenſum 453
— — hirſutulum 439
— — inuitum 443
— — latifolium 457
— — lunulatum 456
— — medullare 452 p. 82
— — nudum 446
— — nymphale 442
— — pennigerum 444
— — puſtulatum 436
— — ſcandens 437
— — ſerpens 435
— — ſetoſum 447
— — tenellum 440
— — veſtitum 445
Polyſcias pinnata 518
Pometia pinnata 392
— ternata 393
Portlandia tetrandra 86
Portulaca lutea 520
Pſidium Decaſpermum 219
Pſychotria aſiatica 90
— — herbacea 91
— — ſpecioſa 89
Pteris comans 419
— eſculenta 418
— humilis 421
— pedata 417
— rotundifolia 420

R

Ranunculus hirtus 525
— — riuularis 524
Reſtis ſimplex 367
Rhamnus polifolius 509
— — Zizyphoides 510
Rhizophora gymnorhiza 201
— — Mangle 202
Rhus atrum 142
Ricinus dioicus 357
— — Mappa 356
Rondeletioides 511
Rottbölla Coelorachis 49
— — repens 50
Rubus auſtralis 224
Ruellia fragrans 243
— — reptans 242
Rumex flexuoſus 515

S

Saccharum officinarum 33
— — ſpontaneum 32
Salicornia auſtralis 489
Salſola Kali 130
— fruticoſa 131
Sapindus Saponaria 178
Scaeuola Lobelia 87
— — ſaligna 505
— — ſericea 504
Schefflera digitata 146
Schoenus arundinaceus 491
— — eleuatus 494
— — rubiginoſus 493
— — ſurinamenſis 492
Scirpus dichotomus 28
— junceus 29
Scorzonera ſcapigera 534
Scrophularioides arborea No. 528
Senecio anguſtifolius 539
— lautus 538
Serapias regularis 312
Seſuvium Portulacaſtrum 222
Shawia paniculata 307
Sheffieldia repens 67
Sicyos angulata 368
Sida 502
— indica 257
— rhombifolia 256
Sigesbeckia orientalis 303
Siſymbrium heterophyllum 250
Siſyrinchium Ixioides 325
Skinnera excorticata 163
Smilax purpurata 373
— Ripogonum 372
Solanum aitroites 508
— — auiculare 107
— — nigrum 106
— — repandum 105
— — viride 507
Solidago arboreſcens 298
Sonchus oleraceus 282
Sophora tomentoſa 182
— — tetraptera 183
Spondias dulcis 198
Stachys decemdentata 526
Sterculia Balanghas 358
— — foetida 359
Suriana maritima 199

T

Tabernaemontana citrifolia 124
Tacca pinnatifida 209
Terminalia Catappa 388
— — glabrata 389

Tetragonia

Tetragonia hallmifolia No. 223
Teucrium villoſum 232
Thalia cannaeformis 3
Tillaea muſcoſa? 61
Tournefortia argentea 64
Trichilia alliacea 189
— ſpectabilis 188
Trichomanes biualue 466
— — contiguum 463
— — cuneiforme 469
— — demiſſum 468
— — demiſſum 467
— — elatum 474
— — epiphyllum 471
— — flaccidum 472
— — gibberoſum 470
— — humile 464
— — multifidum 473
— — reniforme 462
— — ſanguinolentum 465
— — ſolidum 475
— — ſquarroſum 476
Triumfetta procumbens 204
Typha latifolia 336

U

Ulva 591
Urena lobata 258
Urtica aeſtuans 342
Urtica argentea No. 343
— ferox 346
— ruderalis 344
— virgata 345

V

Vaccinium cereum 167
Verbena nodiflora 17
Veronica Catarractae 9
— — elliptica 10
— — ſalicifolia 11
Viſcum antarcticum 370
— opuntioides 369
Vitex trifolia 245
Volkameria inermis 244

W

Waltheria indica 251
— — Lopanthus 252
Weinmannia racemoſa 173
— — paruiflora 174
Wintera axillaris 229

X

Xeranthemum Bellidioides 293
Ximenia elliptica 162
Xylophylla longifolia 143
Xyloſma orbiculatum 381
— — ſuaueolens 380

3 2044 103 107 8

U. HOLZER
BINDER
BOSTON, MASS.